Lecture notes on Free Boundary Problems for Parabolic Equations

DOCTORAL SCHOOL OF CISTERNA, MAY 2011

Daniele Andreucci Dipartimento di Scienze di Base e Applicate per l'Ingegneria Sapienza Università di Roma via Antonio Scarpa 16 00161 Roma, Italy andreucci@dmmm.uniroma1.it

stefan s
dc 2011051610.21

Introduction

These Lecture Notes cover the course Problemi a frontiera libera per equazioni paraboliche Free boundary problems for parabolic equations held in the framework of the Scuola Dottorale di Cisterna (Latina) from 9th to 13th May 2011. The present notes assume the reader has some knowledge of the elementary theory of L^p and Sobolev spaces, as well as of the basic results of existence and regularity of solutions to smooth parabolic equations. The bibliography is minimal; only books and articles quoted in the text are referenced. See [12], [18] and [20] for further references. The present version corrects and supersedes the preliminary version released before the course was completed.

Rome, May 2011

Notation

The notation employed here is essentially standard. Appendix E contains a list of the main symbols used in the text. Constants denoted by γ , C, C_{δ} , ... may change fom line to line.

Contents

Introduction Notation	iii iii
 Chapter 1. The classical formulation 1.1. The Stefan condition 1.2. The free boundary problem 1.3. Explicit examples of solutions 1.4. Basic estimates 1.5. Existence and uniqueness of the solution 1.6. Qualitative behaviour of the solution 1.7. Regularity of the free boundary 	$egin{array}{c} 1 \\ 1 \\ 4 \\ 6 \\ 7 \\ 9 \\ 14 \\ 16 \end{array}$
 Chapter 2. Weak formulation of the Stefan problem 2.1. From the energy balance to the weak formulation 2.2. Definition of weak solution 2.3. Comparing the weak and the classical formulations 2.4. Uniqueness of the weak solution 2.5. Existence of weak solutions 2.6. A comparison result 	19 19 20 21 25 30 38
Chapter 3. The relaxed formulation3.1. A relaxed Stefan problem3.2. Existence of solutions to the relaxed problem3.3. The Stefan problem as limit of the relaxed problem	$ 41 \\ 41 \\ 42 \\ 45 $
 Appendix A. Maximum principles for parabolic equations A.1. The weak maximum principle A.2. The strong maximum principle A.3. Hopf's lemma (the boundary point principle) A.4. Maximum principle for weak solutions 	49 49 51 55 59
Appendix B. A theorem by KruzhkovB.1. Mollifying kernelsB.2. The main estimate	
Appendix C. The spaces $H^{m+\alpha,\frac{m+\alpha}{2}}(\overline{Q_T})$ C.1. Comments	$\begin{array}{c} 65 \\ 65 \end{array}$
Appendix D. Solution of a simple variational inequalityD.1. A variational inequalityD.2. Existence of solutions to (P)D.3. Variational formulation	67 67 68 71

Appendix E.	Symbols used in text	75
Bibliography		77

CHAPTER 1

The classical formulation

In this chapter we consider the formulation of the Stefan problem as a classical initial boundary value problem for a parabolic partial differential equation. A portion of the boundary of the domain is a priori unknown (the *free boundary*), and therefore two boundary conditions must be prescribed on it, instead than only one, to obtain a well posed problem.

1.1. The Stefan condition

The Stefan problem ([17]) is probably the simplest mathematical model of a phenomenon of change of phase. When a change of phase takes place, a *latent heat* is either absorbed or released, while the temperature of the material changing its phase remains constant. In the following we denote by L > 0 the latent heat per unit of volume (p.u.v.), and neglect for the sake of simplicity any volume change in the material undergoing the change of phase. We also assume the critical temperature of change of phase to be a constant, θ_0 .

To be specific, consider at time $t = t_0$ a domain A divided by the plane $x_1 = s_0$ into two subdomains. At time $t = t_0$ the sub-domain $A_1 = A \cap \{x_1 < s_0\}$ is filled by water, while $A_2 = A \cap \{x_1 > s_0\}$ is filled by ice. In the terminology of problems of change of phase, A_1 is the *liquid phase* and A_2 is the solid phase. The surface separating the two phases is referred to as the *interface*. Assume also the setting is plane symmetric, that is the temperature θ is a function of x_1 only, besides the time t, and the interface is a plane at all times. Denote by $x_1 = s(t)$ the position of the interface at time t. Note that, due to the natural assumption that temperature is continuous,

$$\theta(s(t)+,t) = \theta(s(t)-,t) = \theta_0, \quad \text{for all } t. \tag{1.1}$$

Assume ice is changing its phase, that is the interface is advancing into the solid phase. Due to the symmetry assumption we stipulate, we may confine ourselves to consider any portion D, say a disk, of the interface at time t_0 . At a later time $t_1 > t_0$ the interface occupies a position $s(t_1) > s(t_0) = s_0$. The cylinder $D \times (s(t_1), s(t_0))$ has been melted over the time interval (t_0, t_1) (see Figure 1.1). The change of phase has therefore absorbed a quantity of heat

volume of the melted cylinder × latent heat p.u.v. = $\operatorname{area}(D)(s(t_1)-s(t_0))L$. (1.2)

The heat must be provided by diffusion, as we assume that no heat source or sink is present. We adopt for heat diffusion Fourier's law

heat flux
$$= -k_i \nabla \theta$$
, (1.3)



FIGURE 1.1. Melting ice

where $k_1 > 0$ is the diffusivity coefficient in water, and $k_2 > 0$ is the diffusivity coefficient in ice (in principle $k_1 \neq k_2$). Thus, the quantity of heat in (1.2) must equal

$$\int_{t_0}^{t_1} \int_{D(t)} \left[-k_1 \nabla \theta(s(t) - t) \cdot \mathbf{e_1} - k_2 \nabla \theta(s(t) + t) \cdot (-\mathbf{e_1}) \right] dx_2 dx_3 dt = \operatorname{area}(D) \int_{t_0}^{t_1} \left[-k_1 \theta_{x_1}(s(t) - t) + k_2 \theta_{x_1}(s(t) + t) \right] dt. \quad (1.4)$$

Equating the two quantities, dividing the equation by $t_1 - t_0$ and letting $t_1 \rightarrow t_0$, we finally find

$$-k_1\theta_{x_1}(s(t), t) + k_2\theta_{x_1}(s(t), t) = L\dot{s}(t), \qquad (1.5)$$

where we have substituted t_0 with the general time t, as the same procedure can be obviously carried out at any time. This is called the *Stefan condition* on the free boundary. We stress the fact that the Stefan condition is merely a law of energetical balance.

Several remarks are in order.

Remark 1.1. Note that, although we did not assume anything on the values of $\theta(x_1, t)$ *inside* each one of the two phases, on physical grounds we should expect

$$\theta \ge \theta_0$$
, in water, i.e., in A_1 ; $\theta \le \theta_0$, in ice, i.e., in A_2 . (1.6)

The equality $\theta \equiv \theta_0$ in either one of the two phases (or in both) can not be ruled out in the model. Rather, it corresponds to the case when a whole phase is at critical temperature. Diffusion of heat, according to (1.3), can not take place in that phase, as $\theta_{x_1} \equiv 0$ there. Thus (1.5) reduces to, e.g., if the solid phase is at constant temperature,

$$-k_1\theta_{x_1}(s(t), t) = L\dot{s}(t).$$
(1.7)

Note that if $\theta > \theta_0$ in water, (1.7) predicts that $\dot{s}(t) > 0$. In other words, melting of ice is predicted by the model, instead of solidification of water. This is consistent with obvious physical considerations.

Problems where one of the two phases is everywhere at the critical temperature are usually referred to (somehow misleadingly) as one phase problems, while the general case where (1.5) is prescribed is the two phases problem. In the latter case, the sign of $\dot{s}(t)$, and therefore the physical behaviour of the system water/ice predicted by the mathematical model, depends on the relative magnitude of the two heat fluxes at the interface. \Box

Remark 1.2. On the interface, which is also known as the free boundary two conditions are therefore prescribed: (1.1) and (1.5).

In the case of a one phase problem, this fact has the following meaningful interpretation in terms of the general theory of parabolic PDE: The boundary of the domain where the heat equation (a parabolic equation based on (1.3)) is posed, contains an a priori unknown portion, corresponding to the interface separating the liquid phase from the solid one. Clearly, if only the Dirichlet boundary condition (1.1) was imposed on it, we could choose *arbitrarily* this part of the boundary, and solve the corresponding initial value boundary problem. Apparently the solution would not, in general, satisfy (1.7). A similar remark applies to solutions found imposing just (1.7) (where now the functional form of the arbitrarily given boundary $x_1 = s(t)$ is explicitly taken into account).

It is therefore evident that on the free boundary both conditions (1.1) and (1.7) should be prescribed in order to have a well posed problem. (Or, anyway, in more general free boundary problems, two different boundary conditions are required.)

Incidentally, this circle of ideas provides the basic ingredient of a possible proof of the existence of solutions: we assign arbitrarily a 'candidate' free boundary s^* and consider the solution θ to the problem, say, corresponding to the data (1.1). Then we define a *transformed* boundary s^{**} exploiting (1.7), i.e.,

$$-k_1\theta_{x_1}(s^*(t)-,t) = L\dot{s}^{**}(t).$$

A fixed point of this transform corresponds to a solution of the complete problem. $\hfill \Box$

Remark 1.3. The meaning of conditions (1.1) and (1.5) in the context of two phases problems is probably better understood in terms of the weak formulation of the Stefan problem, which is discussed below in Chapter 2. We remark here that, actually, one phase problems are just two phases problems with one phase at constant temperature, so that the discussion in Chapter 2 applies to them too.

Remark 1.4. Problems where the temperature restriction (1.6) is not fulfilled, are sometimes called *undercooled* Stefan problems. We do not treat them here, albeit their mathematical and physical interest (see however Subsection 2.3.4 of Chapter 2); let us only recall that they are, in some sense, ill posed.

We also remark that the relaxed Stefan problem considered in Chapter 3 allows undercooling. $\hfill \Box$

DANIELE ANDREUCCI

1.1.1. Exercises.

Exercise 1.5. Write the analogs of Stefan condition (1.5) in the cases of cylindrical and spherical symmetry in \mathbb{R}^3 .

Exercise 1.6. Note that if we assume, in (1.5), $\theta < \theta_0$ in A_1 and $\theta \equiv \theta_0$ in A_2 , we find $\dot{s}(t) < 0$. This appears to be inconsistent with physical intuition: a phase of ice at sub critical temperature should *grow* into a phase of water at identically critical temperature. Indeed, (1.5) is not a suitable model for the physical setting considered here. In other words, the Stefan condition in the form given above keeps memory of which side of the interface is occupied by which phase. Find out where we implicitly took into account this piece of information and write the Stefan condition when ice and water switch places. \square

Exercise 1.7. Prove that Stefan condition (1.5) does not change its form if bounded volumetric heat sources are present (i.e., if the heat equation is not homogeneous). \square

1.2. The free boundary problem

Keeping the plane symmetry setting considered above, we may of course assume the problem is one dimensional. Denoting by x the space variable, the complete two phases problem can be written as

$$c_1\theta_t - k_1\theta_{xx} = 0, \qquad \text{in } Q_1, \qquad (1.8)$$

$$\begin{aligned} & e_2\theta_t - k_2\theta_{xx} = 0, & \text{in } Q_2, \\ & -k_1\theta_x(0,t) = h_1(t), & 0 < t < T, \end{aligned}$$
(1.9)

$$-\kappa_1 \theta_x(0,t) = h_1(t), \qquad 0 < t < T, \qquad (1.10)$$
$$-\kappa_2 \theta_x(d,t) = h_2(t), \qquad 0 < t < T, \qquad (1.11)$$

$$\theta(x,0) = \Theta(x), \qquad 0 < x < d, \qquad (1.12)$$

$$-k_1 \theta_x(s(t) - t) + k_2 \theta_x(s(t) + t) = L\dot{s}(t), \qquad 0 < t < T, \qquad (1.13)$$

$$\theta(s(t) - t) = \theta(s(t) + t) = \theta_0, \qquad 0 < t < T, \qquad (1.14)$$

$$s(0) = b$$
. (1.15)

 $(1 \ 1 \ 0)$

Here 0 < b < d, T and c_1 , c_2 , k_1 , k_2 are given positive numbers. The c_i represent the thermal capacities in the two phases. The liquid phase occupies at the initial time t = 0 the interval (0, b), while the solid phase occupies (b, d). The problem is posed in the time interval (0, T). Moreover we have set

$$Q_1 = \{(x,t) \mid 0 < x < s(t), \ 0 < t < T\},\$$
$$Q_2 = \{(x,t) \mid s(t) < x < d, \ 0 < t < T\}.$$

We are assuming that 0 < s(t) < d for all 0 < t < T. If the free boundary hits one of the two fixed boundaries x = 0 and x = d, say at time t^* , of course the formulation above should be changed. In practice, one of the two phases disappears at $t = t^*$. We leave to the reader the simple task of writing the mathematical model for $t > t^*$.

One could impose other types of boundary data, instead of (1.10), (1.11), e.g., Dirichlet data.

If we are to attach the physical meaning of a change of phase model to the problem above, the data must satisfy suitable conditions. At any rate

 $\Theta(x) \ge \theta_0$, 0 < x < b; $\Theta(x) \le \theta_0$, b < x < d.

Essentially, we need $\theta \geq \theta_0$ in Q_1 and $\theta \leq \theta_0$ in Q_2 .

Actually, we will deal mainly with the one phase version of (1.8)-(1.15)where the solid phase is at constant temperature. Namely, after adimensionalization, we look at

$$u_t - u_{xx} = 0,$$
 in $Q_{s,T},$ (1.16)

$$-u_x(0,t) = h(t) > 0, \qquad 0 < t < T, \qquad (1.17)$$
$$u(x,0) = u_0(x) > 0, \qquad 0 < x < b. \qquad (1.18)$$

$$u(x,0) = u_0(x) \ge 0, \qquad 0 < x < b,$$
 (1.18)

$$u_x(s(t), t) = \dot{s}(t), \qquad 0 < t < T, \qquad (1.10)$$

$$u(s(t), t) = 0,$$
 $0 < t < T,$ (1.20)

$$s(0) = b$$
. (1.21)

(We have kept the old names for all variables excepting the unknown u.) Here we denote for each positive function $s \in C([0, T])$, such s(0) = b,

$$Q_{s,T} = \{(x,t) \mid 0 < x < s(t), 0 < t < T\}.$$

We regard the rescaled temperature u as a function defined in $Q_{s,T}$. The solid phase therefore does not appear explicitly in the problem. As a matter of fact we assume it to be unbounded in the positive x direction (i.e., d = $+\infty$), so that no upper limit has to be imposed on the growth of the free boundary s. The sign restrictions in (1.17) and in (1.18) are imposed so that u > 0 in $Q_{s,T}$, see Proposition 1.12 below.

Definition 1.8. A solution to problem (1.16)-(1.21) is a pair (u, s) with

$$s \in C^{1}((0,T]) \cap C([0,T]), \quad s(0) = b, \quad s(t) > 0, \ 0 \le t \le T;$$
$$u \in C(\overline{Q_{s,T}}) \cap C^{2,1}(Q_{s,T}), \qquad u_{x} \in C(\overline{Q_{s,T}} - \{t = 0\}),$$

and such that (1.16)-(1.21) are satisfied in a classical pointwise sense.

We prove a theorem of existence and uniqueness of solutions to (1.16)-(1.21), under the assumptions

$$h \in C([0,T]), \qquad h(t) > 0, \ 0 \le t \le T;$$
 (1.22)

$$u_0 \in C([0,b]), \qquad 0 \le u_0(x) \le H(b-x), \ 0 \le x \le b.$$
 (1.23)

We also study some qualitative behaviour of the solution. The adimensionalization of the problem does not play a substantial role in the mathematical theory we develop here.

For further reading on the one-phase Stefan problem, we refer the reader to [5], [3]; we employ in this chapter the techniques found there, with some changes.

Remark 1.9. The free boundary problem (1.16)-(1.21) is strongly non linear, in spite of the linearity of the PDE and of the boundary conditions there. Indeed, recall that the free boundary s itself is an unknown of the problem; its dependence on the data is not linear (as, e.g., the explicit examples of Section 1.3 show). \square

DANIELE ANDREUCCI

1.2.1. Exercises.

Exercise 1.10. Prove that the change of variables

$$\theta \mapsto \alpha u + \theta_0, \quad x \mapsto \beta \xi, \quad t \mapsto \gamma \tau$$

allows one to write the one phase problem in the adimensionalized form (1.16)-(1.21), for a suitable choice of the constants α , β , γ . Also note that adimensionalizing the complete two phases problem similarly

Also note that admensionalizing the complete two phases problem similarly is in general impossible. \Box

1.3. Explicit examples of solutions

EXAMPLE 1. An explicit solution of the heat equation (1.16) is given by

$$v(x,t) = \operatorname{erf}\left(\frac{x}{2\sqrt{t}}\right) = \frac{2}{\sqrt{\pi}} \int_{0}^{x/2\sqrt{t}} e^{-z^{2}} dz, \qquad x, t > 0,$$
$$v(x,0) = 1, \qquad x > 0.$$

Here erf denotes the well known 'error function'. Fix C > 0 arbitrarily, and set, for a $\alpha > 0$ to be chosen presently,

$$u(x,t) = C\left\{\operatorname{erf} \alpha - \operatorname{erf}\left(\frac{x}{2\sqrt{t}}\right)\right\}.$$

Define also $s(t) = 2\alpha\sqrt{t}$; note that s(0) = 0. Thus u > 0 in $Q_{s,T}$, and (1.16) as well as (1.20) are satisfied. By direct calculation

$$u_x(x,t) = -\frac{C}{\sqrt{\pi t}}e^{-\frac{x^2}{4t}}.$$

Hence,

$$u_x(s(t),t) = u_x(2\alpha\sqrt{t},t) = -\frac{C}{\sqrt{\pi t}}e^{-\alpha^2} = -\dot{s}(t) = -\frac{\alpha}{\sqrt{t}},$$

if and only if

$$C = \sqrt{\pi} \alpha e^{\alpha^2} \,.$$

Note that on the fixed boundary x = 0 we may select either one of the conditions

$$u_x(0,t) = -\frac{C}{\sqrt{\pi t}}$$
, or $u(0,t) = C \operatorname{erf} \alpha$.

We have proven that, when α is chosen as above,

$$u(x,t) = 2\alpha e^{\alpha^2} \int_{x/2\sqrt{t}}^{\alpha} e^{-z^2} dz \qquad (1.24)$$

solves the problem sketched in Figure 1.2. Note however that u is not continuous at (0,0); the notion of solution in this connection should be suitably redefined.

EXAMPLE 2. It is obvious by direct inspection that the function

$$u(x,t) = e^{t-x} - 1, \qquad (1.25)$$



FIGURE 1.2. The Stefan problem solved by u in (1.24)

solves the Stefan problem in Figure 1.3, corresponding to the free boundary s(t) = t. Note that we are forced to prescribe an exponentially increasing flux on x = 0 in order to obtain a linear growth for s(t).



FIGURE 1.3. The Stefan problem solved by u in (1.25)

1.3.1. Exercises.

Exercise 1.11. Convince yourself that the solutions corresponding to $u_x(0,t) = -2C/\sqrt{\pi t}$, in the case of Example 1, and to $u_x(0,t) = -2e^t$ in the case of Example 2, can not be obtained by linearity from the ones given above. \Box

1.4. Basic estimates

Proposition 1.12. If (u, s) is a solution to (1.16)–(1.21), then

$$u(x,t) > 0, \qquad in \ Q_{s,T};$$
 (1.26)

$$\dot{s}(t) > 0$$
, for all $t > 0$. (1.27)

PROOF. By virtue of the weak maximum principle, u must attain its minimum on the parabolic boundary of $Q_{s,T}$, i.e., on

$$\partial Q_{s,T} - \{(x,t) \mid t = T, 0 < x < s(T)\}.$$

DANIELE ANDREUCCI

The data h being positive, the minimum is attained on t = 0 or on x = s(t). Therefore $u \ge 0$ (remember that $u_0 \ge 0$). If we had $u(\bar{x}, \bar{t}) = 0$ in some $(\bar{x}, \bar{t}) \in Q_{s,T}$, invoking the strong maximum principle we would obtain $u \equiv 0$ in $Q_{s,T} \cap \{t < \bar{t}\}$. This is again inconsistent with h > 0. Thus (1.26) is proven.

Then, the value u = 0 attained on the free boundary is a minimum for u. Recalling the parabolic version of Hopf's lemma, we infer

$$\dot{s}(t) = -u_x(s(t), t) > 0$$
, for all $t > 0$.

Remark 1.13. The proof of estimate (1.26) does not make use of the Stefan condition (1.19). In the same spirit, we consider in the following results solutions to the initial value boundary problem obtained removing Stefan condition from the formulation. The rationale for this approach is that we want to apply those results to 'approximating' solutions constructed according to the ideas of Remark 1.2.

Lemma 1.14. Let u be a solution of (1.16), (1.17), (1.18), (1.20), where s is assumed to be a positive non decreasing Lipschitz continuous function in [0,T], such that s(0) = b. Let (1.22) and (1.23) be in force. Then u_x is continuous up to all the points of the boundary of $Q_{s,T}$ of the form (0,t), (s(t),t), with $T \ge t > 0$.

The regularity of u_x up to x = 0 is classical; the proof of this result will be completed in Section 1.7, see Lemma 1.31.

Proposition 1.15. Let u, s be as in Lemma 1.14. Then

$$0 < u(x,t) \le M(s(t) - x), \qquad in \ Q_{s,T}, \tag{1.28}$$

where $M = \max(\|h\|_{\infty}, H)$.

v

PROOF. Define v(x,t) = M(s(t) - x). It follows immediately

$$v_t - v_{xx} = M\dot{s}(t) \ge 0, \quad \text{in } Q_{s,T};$$

$$v(s(t),t) = u(s(t),t) = 0, \quad 0 \le t \le T,$$

$$v_x(0,t) = -M \le -h = u_x(0,t), \quad 0 < t < T,$$

$$(x,0) = M(b-x) \ge u_0(x) = u(x,0), \quad 0 \le x \le b.$$

Therefore, taking into account the results of Appendix A,

$$v(x,t) \ge u(x,t), \quad \text{in } Q_{s,T}.$$

Corollary 1.16. If u, s are as in Proposition 1.15, then

$$0 > u_x(s(t), t) \ge -M, \qquad 0 < t < T, \tag{1.29}$$

where M is the constant defined in Proposition 1.15.

PROOF. A trivial consequence of Proposition 1.12 and of Proposition 1.15, as well as of Lemma 1.14. We also keep in mind Remark 1.13, and of course make use of Hopf's lemma for the strict inequality in (1.29).

Remark 1.17. We stress the fact that the barrier construction of Proposition 1.15 is made possible by the fact that $\dot{s} \ge 0$. In turn, this is for solutions of the Stefan problem a consequence of the positivity of u. Thus, such a barrier function, and in general any similar barrier function, does not exist in the case of the undercooled Stefan problem.

1.4.1. Exercises.

Exercise 1.18. Note that the function v defined in the proof of Proposition 1.15 is just Lipschitz continuous in t. In spite of this fact, one may apply to v - u the weak maximum principle in the form given in Section A.4 of Appendix A. Carry out the proof in detail.

1.5. Existence and uniqueness of the solution

We prove here

Theorem 1.19. Assume (1.22), (1.23). Then there exists a unique solution to (1.16)–(1.21).

Let u, s be as in Proposition 1.15. Moreover assume that

$$s \in \Sigma := \{ \sigma \in \operatorname{Lip}([0,T]) \mid 0 \le \dot{\sigma} \le M, \, \sigma(0) = b \}$$

Here M is the constant defined in Proposition 1.15. The set Σ is a convex compact subset of the Banach space C([0, T]), equipped with the max norm. A useful property of all $s \in \Sigma$ is

$$b + Mt \ge s(t) \ge b$$
, $0 \le t \le T$.

Define the transform $\mathcal{T}(s)$ by

$$\mathcal{T}(s)(t) = b - \int_0^t u_x(s(\tau), \tau) \,\mathrm{d}\tau, \qquad T \ge t \ge 0.$$

Note that, as a consequence of Lemma 1.14 and of Corollary 1.16,

$$\mathcal{T}(s) \in \operatorname{Lip}([0,T]) \cap C^1((0,T]),$$

and

$$M \ge \frac{\mathrm{d}}{\mathrm{d}t} \mathcal{T}(s)(t) = -u_x(s(t), t) \ge 0, \qquad t > 0.$$

Then $\mathcal{T} : \Sigma \to \Sigma$. Also note that a fixed point of \mathcal{T} corresponds to a solution of our Stefan problem.

We use the divergence theorem to transform the boundary flux integral defining $\mathcal{T}(s)$. Namely we write

$$0 = \int_{0}^{t} \int_{0}^{s(\tau)} (u_{\tau} - u_{xx}) \, \mathrm{d}x \, \mathrm{d}\tau = -\int_{0}^{b} u_{0}(x) \, \mathrm{d}x - \int_{0}^{t} u(s(\tau), \tau) \dot{s}(\tau) \, \mathrm{d}\tau + \int_{0}^{s(t)} u(x, t) \, \mathrm{d}x - \int_{0}^{t} u_{x}(s(\tau), \tau) \, \mathrm{d}\tau - \int_{0}^{t} h(\tau) \, \mathrm{d}\tau$$

Therefore

$$\mathcal{T}(s)(t) = b - \int_{0}^{t} u_{x}(s(\tau), \tau) \,\mathrm{d}\tau =$$

$$b + \int_{0}^{b} u_{0}(x) \,\mathrm{d}x + \int_{0}^{t} h(\tau) \,\mathrm{d}\tau - \int_{0}^{s(t)} u(x, t) \,\mathrm{d}x =: F(t) - \int_{0}^{s(t)} u(x, t) \,\mathrm{d}x. \quad (1.30)$$

Note that this equality allows us to express $\mathcal{T}(s)$ in terms of more regular functions than the flux $u_x(s(t), t)$, which appeared in its original definition. We are now in a position to prove that \mathcal{T} is continuous in the max norm. Let $s_1, s_2 \in \Sigma$. Let us define

$$\alpha(t) = \min(s_1(t), s_2(t)), \qquad \beta(t) = \max(s_1(t), s_2(t)),$$

 $i = 1, \quad \text{if } \beta(t) = s_1(t), \qquad i = 2, \quad \text{otherwise.}$

(The number i is a function of time; this will not have any specific relevance.) Let us also define

$$v(x,t) = u_1(x,t) - u_2(x,t)$$

Then v satisfies

$$v_t - v_{xx} = 0,$$
 in $Q_{\alpha,T},$ (1.31)

$$v_x(0,t) = 0,$$
 $0 < t < T,$ (1.32)

$$v(x,0) = 0, \qquad 0 < x < b, \qquad (1.33)$$

$$|v(\alpha(t), t)| = |u_i(\alpha(t), t)| \le M(\beta(t) - \alpha(t)), \qquad 0 < t < T.$$
(1.34)

Therefore, we may invoke the maximum principle to obtain

$$\|v\|_{\infty,t} := \max_{\overline{Q}_{\alpha,t}} |v| \le M \|s_1 - s_2\|_{\infty,t}, \qquad (1.35)$$

where we also denote

$$||s_1 - s_2||_{\infty,t} = \max_{0 \le \tau \le t} |s_1(\tau) - s_2(\tau)|.$$

On the other hand, we have

$$\mathcal{T}(s_1)(t) - \mathcal{T}(s_2)(t) = \int_{0}^{s_2(t)} u_2(x,t) \, \mathrm{d}x - \int_{0}^{s_1(t)} u_1(x,t) \, \mathrm{d}x$$
$$= -\int_{0}^{\alpha(t)} v(x,t) \, \mathrm{d}x + (-1)^i \int_{\alpha(t)}^{\beta(t)} u_i(x,t) \, \mathrm{d}x$$

Therefore

$$\begin{aligned} |\mathcal{T}(s_1)(t) - \mathcal{T}(s_2)(t)| &\leq |\alpha(t)| \|v\|_{\infty,t} + M(\beta(t) - \alpha(t))^2 \\ &\leq (b + MT)M \|s_1 - s_2\|_{\infty,t} + M \|s_1 - s_2\|_{\infty,t}^2, \quad (1.36) \end{aligned}$$

and the continuity of $\mathcal{T} : \Sigma \to \Sigma$ is proven. By Schauder's theorem, it follows that a fixed point of \mathcal{T} exists, and thus existence of a solution.

Uniqueness might be proven invoking the monotone dependence result given below (see Theorem 1.27).

However, mainly with the purpose of elucidating the role played in the theory of free boundary problems by local integral estimates, we proceed to give a direct proof of uniqueness of solutions. More explicitly, we prove the contractive character (for small t) of \mathcal{T} , thereby obtaining existence and uniqueness of a fixed point.

1.5.1. Local estimates vs the maximum principle. Estimates like (1.35), obtained through the maximum principle, have the advantage of providing an immediate sup estimate of the solution in the whole domain of definition. However, the bound they give may be too rough, at least in some regions of the domain. Consider for example, in the setting above, a point $(b/2, \varepsilon)$, with $\varepsilon \ll 1$. The maximum principle predicts for $v(b/2, \varepsilon)$ a bound of order $M ||s_1 - s_2||_{\infty,\varepsilon}$, that is, obviously, the same bound satisfied by the boundary data for v. On the other hand, taking into account (1.32), (1.33) one might expect $v(b/2, \varepsilon)$ to be much smaller than $v(\alpha(\varepsilon), \varepsilon)$.

This is indeed the case, as we show below. In order to do so, we exploit local integral estimates of the solution, that is, estimates involving only values of v in the region of interest, in this case, away from the boundary. We aim at proving that \mathcal{T} is a contraction, that is,

$$\|\mathcal{T}(s_1) - \mathcal{T}(s_2)\|_{\infty,t} \le d\|s_1 - s_2\|_{\infty,t}, \qquad (1.37)$$

with d < 1 for small enough t. A quick glance at (1.36) shows that (1.37) does not, indeed, follow from there (unless bM < 1). This failure is due only to the term originating from the estimate of

$$\int_{0}^{\alpha(t)} v(x,t) \,\mathrm{d}x$$

Thus, a better estimate of this integral is needed.

A key step in any local estimation is a good choice of cut off functions. These are, typically, non negative smooth functions equal to 1 in the region we want to single out, and identically vanishing away from it.

Let $b/2 > \delta > 0$, and define the cut off function $\zeta(x)$, such that

$$\zeta(x) = 1$$
, $0 \le x \le b - 2\delta$, $\zeta(x) = 0$, $b - \delta \le x$, $-\frac{2}{\delta} \le \zeta_x(x) \le 0$.

Multiply (1.31) by $v\zeta^2$ and integrate by parts in $Q_{b,t} = [0,b] \times [0,t]$. We get

$$0 = \iint_{Q_{b,t}} (v_{\tau} v \zeta^{2} - v_{xx} v \zeta^{2}) \, \mathrm{d}x \, \mathrm{d}\tau$$

= $\frac{1}{2} \int_{0}^{b} v(x,t)^{2} \zeta(x)^{2} \, \mathrm{d}x + \iint_{Q_{b,t}} v_{x}^{2} \zeta^{2} \, \mathrm{d}x \, \mathrm{d}\tau + 2 \iint_{Q_{b,t}} v v_{x} \zeta \zeta_{x} \, \mathrm{d}x \, \mathrm{d}\tau \,, \quad (1.38)$

whence (using the inequality $2ab \leq \varepsilon a^2 + b^2/\varepsilon, \varepsilon > 0$)

$$\frac{1}{2} \int_{0}^{b} v(x,t)^{2} \zeta(x)^{2} \,\mathrm{d}x + \iint_{Q_{b,t}} v_{x}^{2} \zeta^{2} \,\mathrm{d}x \,\mathrm{d}\tau \leq 2 \iint_{Q_{b,t}} |v_{x}| \,|v| \,|\zeta_{x}|\zeta \,\mathrm{d}x \,\mathrm{d}\tau$$
$$\leq 2 \iint_{Q_{b,t}} v^{2} \zeta_{x}^{2} \,\mathrm{d}x \,\mathrm{d}\tau + \frac{1}{2} \iint_{Q_{b,t}} v_{x}^{2} \zeta^{2} \,\mathrm{d}x \,\mathrm{d}\tau + \frac{1}{2} \iint_{Q_{b,t}} v_{x}^{2} \zeta^{2} \,\mathrm{d}x \,\mathrm{d}\tau$$

By absorbing the last integral into the left hand side, we find

$$\int_{0}^{b} v(x,t)^{2} \zeta(x)^{2} dx + \iint_{Q_{b,t}} v_{x}^{2} \zeta^{2} dx d\tau \leq 4 \iint_{Q_{b,t}} v^{2} \zeta_{x}^{2} dx d\tau$$
$$= 4 \int_{0}^{t} \int_{b-2\delta}^{b-\delta} v^{2} \zeta_{x}^{2} dx d\tau \leq \frac{16}{\delta^{2}} t \delta ||v||_{\infty,t}^{2} = \frac{16}{\delta} t ||v||_{\infty,t}^{2} . \quad (1.39)$$

Let us now go back to

$$\mathcal{T}(s_1)(t) - \mathcal{T}(s_2)(t) = -\int_{0}^{\alpha(t)} v(x,t) \, \mathrm{d}x + (-1)^i \int_{\alpha(t)}^{\beta(t)} u_i(x,t) \, \mathrm{d}x$$
$$= \int_{0}^{b-2\delta} v(x,t) \, \mathrm{d}x + \int_{b-2\delta}^{\alpha(t)} v(x,t) \, \mathrm{d}x + (-1)^i \int_{\alpha(t)}^{\beta(t)} u_i(x,t) \, \mathrm{d}x.$$

Use Hölder's inequality and (1.39) to bound the integral over $(0, b - 2\delta)$ (recall that $\zeta \equiv 1$ there), and find

$$|\mathcal{T}(s_1)(t) - \mathcal{T}(s_2)(t)| \le \left(\int_{0}^{b-2\delta} v(x,t)^2 \,\mathrm{d}x\right)^{1/2} \sqrt{b} + (\alpha(t) - b + 2\delta) \|v\|_{\infty,t} + M(\beta(t) - \alpha(t))^2 \le \sqrt{b} \frac{4}{\sqrt{\delta}} \sqrt{t} \|v\|_{\infty,t} + (Mt + 2\delta) \|v\|_{\infty,t} + M(\beta(t) - \alpha(t))^2.$$

Take now $\delta = \sqrt{t}$ (t is fixed in this argument), and apply again the sup estimate (1.35), finally obtaining

$$\|\mathcal{T}(s_1) - \mathcal{T}(s_2)\|_{\infty,t} \le \left\{ 4M\sqrt{b}\sqrt[4]{t} + M(Mt + 2\sqrt{t}) + M^2t \right\} \|s_1 - s_2\|_{\infty,t}.$$

Clearly, for $t = t_0(M, b)$, \mathcal{T} is a contractive mapping.

1.5.2. Exercises.

Exercise 1.20. In Subsection 1.5.1 we proved existence and uniqueness of a fixed point of \mathcal{T} in a small time interval (*locally in time*). Show how the argument can be completed to give existence and uniqueness of a fixed point in any time interval (*global existence*).

Exercise 1.21. Why uniqueness of solutions (u, s) with $s \in \Sigma$ is equivalent to uniqueness of solutions in the class of Definition 1.8, without further restrictions?

Exercise 1.22. Give an interpretation of (1.30) as an energetical balance.

Exercise 1.23. Prove that \mathcal{T} has the property (see also Figure 1.4)

$$s_1(t) < s_2(t), \quad 0 < t < T \implies \mathcal{T}(s_1)(t) > \mathcal{T}(s_2)(t), \quad 0 < t < T.$$

Exercise 1.24. Using Exercise 1.23 give an alternative proof of the existence of the fixed point of \mathcal{T} by the following argument. Define

 $s_1(t) = b + Mt$, $s_{n+1} = \mathcal{T}(s_n)$, for $n \ge 1$.

Show that the following monotonic limits exist and are uniform in \boldsymbol{t}

$$s_{2k+1} \searrow \sigma_1, \quad s_{2k} \nearrow \sigma_2, \quad \sigma_1 \ge \sigma_2.$$

Notice that

$$\mathcal{T}(\sigma_1) = \sigma_2, \qquad \mathcal{T}(\sigma_2) = \sigma_1$$

Finally, $\sigma_1 = \sigma_2$ can be proven by the arguments in Subsection 1.5.1.

Exercise 1.25. Extend the proof of existence and uniqueness of solutions to the case when $h \ge 0$. What happens if $h \equiv 0$, $u_0 \equiv 0$?

Exercise 1.26. To carry out rigorously the calculations in (1.38) actually we need an approximation procedure: i.e., we need first perform integration in a smaller 2-dimensional domain, bounded away from the boundaries x = 0, t = 0. Recognize the need of this approach, and go over the (easy) details.



FIGURE 1.4. Behaviour of the transform \mathcal{T}

DANIELE ANDREUCCI

1.6. Qualitative behaviour of the solution

Theorem 1.27. (MONOTONE DEPENDENCE) Let (u_i, s_i) be solutions of (1.16)-(1.21), i = 1, 2, respectively corresponding to data $h = h_i$, $b = b_i$, $u_0 = u_{0i}$. Assume both sets of data satisfy (1.22), (1.23). If

 $h_1(t) \le h_2(t)$, 0 < t < T; $b_1 \le b_2$; $u_{01}(x) \le u_{02}(x)$, $0 < x < b_1$; (1.40)

then

$$s_1(t) \le s_2(t), \qquad 0 < t < T.$$
 (1.41)

PROOF. 1) Let us assume first $b_1 < b_2$. Reasoning by contradiction, assume

$$\overline{t} = \inf\{t \mid s_1(t) = s_2(t)\} \in (0, T).$$

Then the function $v = u_2 - u_1$ is strictly positive in

$$\{0 < t < \overline{t}, 0 < x < s_1(t)\},\$$

by virtue of the strong maximum principle and Hopf's lemma. Indeed,

$$v(s_1(t), t) > 0, \qquad 0 < t < \bar{t}.$$

Then v attains a minimum at $(s_1(\bar{t}), \bar{t}) = (s_2(\bar{t}), \bar{t})$, where

$$v(s_1(\bar{t}),\bar{t})=0.$$

Thus, due to Hopf's lemma,

$$v_x(s_1(\bar{t}),\bar{t}) < 0.$$

But we compute

$$v_x(s_1(\bar{t}), \bar{t}) = u_{2x}(s_2(\bar{t}), \bar{t}) - u_{1x}(s_1(\bar{t}), \bar{t}) = -\dot{s}_2(\bar{t}) + \dot{s}_1(\bar{t}).$$

Hence $\dot{s}_2(\bar{t}) > \dot{s}_1(\bar{t})$, which is not consistent with the definition of \bar{t} . 2) Assume now $b_1 = b_2$. Let us extend the data u_{02} to zero over $(b, b + \delta)$, where $0 < \delta < 1$ is arbitrary. Let (u_{δ}, s_{δ}) be the solution of problem (1.16)–(1.21) corresponding to the data $h = h_2$, $b = b_2 + \delta$, $u_0 = u_{02}$. Then, by the first part of the proof, $s_2 < s_{\delta}$, $s_1 < s_{\delta}$, and for all 0 < t < T

$$s_{\delta}(t) - s_{2}(t) = \delta - \int_{0}^{s_{2}(t)} [u_{\delta} - u_{2}](x, t) \, \mathrm{d}x - \int_{s_{2}(t)}^{s_{\delta}(t)} u_{\delta}(x, t) \, \mathrm{d}x \le \delta \,.$$

Therefore $s_{\delta} \leq s_2 + \delta$, so that $s_1 < s_{\delta} \leq s_2 + \delta$. On letting $\delta \to 0$ we recover $s_1 \leq s_2$.

Let us investigate the behaviour of the solution of the Stefan problem for large times. In doing so, we of course assume that $T = \infty$. Note that the result of existence and uniqueness applies over each finite time interval; a standard extension technique allows us to prove existence and uniqueness of a solution defined for all positive times. Owing to Proposition 1.15, we have

 $s(t) \le S$, $0 < t < \infty \implies u(x,t) \le MS$, in $Q_{s,\infty}$. (1.42)

Moreover, s being monotonic, certainly there exists

$$s_{\infty} = \lim_{t \to \infty} s(t) \,. \tag{1.43}$$

Theorem 1.28. Let (u, s) be the solution of Theorem 1.19. Then

$$s_{\infty} = \lim_{t \to \infty} s(t) = b + \int_{0}^{b} u_{0}(x) \,\mathrm{d}x + \int_{0}^{\infty} h(t) \,\mathrm{d}t \,. \tag{1.44}$$

PROOF. 1) Assume first

$$\int_{0}^{\infty} h(t) \,\mathrm{d}t = +\infty \,. \tag{1.45}$$

We only need show s is unbounded. Let us recall that, for all t > 0,

$$s(t) = b + \int_{0}^{b} u_{0}(x) \,\mathrm{d}x + \int_{0}^{t} h(\tau) \,\mathrm{d}\tau - \int_{0}^{s(t)} u(x,t) \,\mathrm{d}x \,. \tag{1.46}$$

From (1.42), it follows that, if s is bounded over $(0, \infty)$, then u is also bounded over $(0, \infty)$. This is clearly inconsistent with (1.46), when we keep in mind (1.45).

2) Assume

$$\int_{0}^{\infty} h(t) \,\mathrm{d}t < +\infty \,. \tag{1.47}$$

The balance law (1.46), together with u > 0, immediately yields

$$s(t) < b + \int_{0}^{b} u_{0}(x) \,\mathrm{d}x + \int_{0}^{\infty} h(t) \,\mathrm{d}t < +\infty.$$
 (1.48)

Then we have $s(t) \to s_{\infty} < \infty$; it is only left to identify s_{∞} as the quantity indicated above. Owing to (1.46) again, we only need show

$$\lim_{t \to \infty} \int_{0}^{s(t)} u(x,t) \, \mathrm{d}x = 0.$$
 (1.49)

On multiplying (1.16) by u and integrating by parts in $Q_{s,t}$, we get

$$\frac{1}{2} \int_{0}^{s(t)} u(x,t)^2 \,\mathrm{d}x + \iint_{Q_{s,t}} u_x^2 \,\mathrm{d}x \,\mathrm{d}\tau = \frac{1}{2} \int_{0}^{b} u_0(x)^2 \,\mathrm{d}x + \int_{0}^{t} u(0,\tau)h(\tau) \,\mathrm{d}\tau \,. \tag{1.50}$$

Recalling (1.42) and (1.48), the last integral in (1.50) is majorised by

$$\int_{0}^{\infty} M s_{\infty} h(\tau) \, \mathrm{d}\tau < \infty \, .$$

Thus a consequence of (1.50) is

$$\iint_{Q_{s,\infty}} u_x(x,t)^2 \,\mathrm{d}x \,\mathrm{d}t < +\infty \,. \tag{1.51}$$

Elementary calculus then shows that

$$\iint_{Q_{s,\infty}} u(x,t)^2 \,\mathrm{d}x \,\mathrm{d}t = \iint_{Q_{s,\infty}} \left[\int_x^{s(t)} u_{\xi}(\xi,t) \,\mathrm{d}\xi \right]^2 \,\mathrm{d}x \,\mathrm{d}t$$
$$\leq s_{\infty} \iint_{Q_{s,\infty}} \int_x^{s(t)} u_{\xi}(\xi,t)^2 \,\mathrm{d}\xi \,\,\mathrm{d}x \,\mathrm{d}t \leq s_{\infty}^2 \iint_{Q_{s,\infty}} u_{\xi}(\xi,t)^2 \,\mathrm{d}\xi \,\mathrm{d}t < \infty \,\mathrm{d}t$$

Then there exists a sequence $\{t_n\}, t_n \to \infty$, such that

$$\int_{0}^{s(t_n)} u(x,t_n)^2 \,\mathrm{d}x \to 0 \,.$$

But the function

$$t \mapsto \int_{0}^{s(t)} u(x,t)^2 \,\mathrm{d}x\,,$$

when we take into account (1.50), is easily seen to have limit as $t \to \infty$, so that this limit is 0. By Hölder's inequality,

$$\int_{0}^{s(t)} u(x,t) \, \mathrm{d}x \le \sqrt{s_{\infty}} \left[\int_{0}^{s(t)} u(x,t)^2 \, \mathrm{d}x \right]^{1/2} \to 0, \qquad t \to \infty,$$

completing the proof of (1.49).

1.6.1. Exercises.

Exercise 1.29. Find conditions ensuring that the inequality in (1.41) is strict.

Exercise 1.30. Discuss the necessity of assumptions (1.22), (1.23) in Theorem 1.27.

1.7. Regularity of the free boundary

The approach in this Section is taken from [16], and provides an example of 'bootstrap' argument, i.e., of an inductive proof where any given smoothness of the solution allows us to prove even more regularity for it. Our first result will become the first step in the induction procedure, and is however required to prove Lemma 1.14.

Lemma 1.31. Let $u \in C^{2,1}(Q_{s,T}) \cap C(\overline{Q_{s,T}})$, where $s \in \text{Lip}([0,T])$, and s(t) > 0 for $0 \le t \le T$. Assume u fulfils

$$u_t - u_{xx} = 0, \qquad in \ Q_{s,T},$$
 (1.52)

$$u(s(t), t) = 0, \qquad 0 \le t \le T.$$
 (1.53)

Then for each small enough $\varepsilon > 0$, u_x is continuous in $\overline{P_{\varepsilon}}$, where $P_{\varepsilon} = \{(x,t) \mid \varepsilon < x < s(t), \varepsilon < t < T\}.$

PROOF. The proof is based on standard local regularity estimates for solutions of parabolic equations. Introduce the following change of variables

$$\begin{cases} y = \frac{x}{s(t)}, \\ \tau = t; \end{cases} \quad \quad v(y,\tau) = u(ys(\tau),\tau). \end{cases}$$

The set $Q_{s,T}$ is mapped onto $R = (0,1) \times (0,T)$, where v solves

$$v_{\tau} - \frac{1}{s(\tau)^2} v_{yy} - \frac{\dot{s}(\tau)}{s(\tau)} y v_y = 0, \quad \text{in } R,$$
(1.54)

$$v(1,\tau) = 0, \qquad 0 \le \tau \le T.$$
 (1.55)

More explicitly, (1.54) is solved a.e. in R, as v is locally a Sobolev function in R. Classical results, see [11] Chapter IV, Section 10, imply that for any fixed $\varepsilon > 0$,

$$v_{\tau}, v_{y}, v_{yy} \in L^{q}((\varepsilon, 1) \times (\varepsilon, T)),$$

for all q > 1. Then we use the embedding Lemma 3.3 of [11] Chapter II, to infer that, for q > 3,

$$v_y \in H^{\alpha, \frac{\alpha}{2}}([\varepsilon, 1] \times [\varepsilon, T]), \qquad (1.56)$$

where $\alpha = 1 - 3/q$ (see also Remark 11.2 of [11], p. 218; the space $H^{\alpha, \frac{\alpha}{2}}$ is defined in Appendix C). Since

$$u_x(x,t) = \frac{1}{s(t)} v_y\left(\frac{x}{s(t)},t\right),$$

the result follows.

Our next result implies that the free boundary in the Stefan problem (1.16)–(1.21) is of class $C^{\infty}(0,T)$.

Theorem 1.32. Assume u and s are as in Lemma 1.31, and moreover

$$u_x(s(t), t) = c\dot{s}(t), \qquad 0 < t < T, \qquad (1.57)$$

where $c \neq 0$ is a given constant. Then $s \in C^{\infty}(0,T)$.

PROOF. For v defined as in the proof of Lemma 1.31, we rewrite (1.57) as

$$\dot{s}(\tau) = \frac{1}{cs(\tau)} v_y(1,\tau), \qquad 0 < \tau < T.$$
 (1.58)

Choose $\alpha \in (0, 1)$. Then (1.56) and (1.58) yield at once

$$\dot{s} \in H^{\alpha,\frac{\alpha}{2}}([\varepsilon,1] \times [\varepsilon,T]), \quad \text{for each fixed } \varepsilon > 0.$$
 (1.59)

Next we make use of the following classical result:

If the coefficients in (1.54) (i.e.,
$$\dot{s}$$
), are of class
 $H^{m+\alpha,\frac{m+\alpha}{2}}([\varepsilon,1] \times [\varepsilon,T])$, then v is of class (1.60)
 $H^{2+m+\alpha,\frac{2+m+\alpha}{2}}([2\varepsilon,1] \times [2\varepsilon,T])$.

Here $m \ge 0$ is any integer. We prove by induction that for all $m \ge 0$

$$\dot{s} \in H^{m+\alpha,\frac{m+\alpha}{2}}([\varepsilon,1] \times [\varepsilon,T]), \quad \text{for each fixed } \varepsilon > 0.$$
 (1.61)

We already know this is the case when m = 0, from (1.59). Assume then (1.61) is in force for a given $m \ge 0$. Then, owing to (1.60),

$$v \in H^{2+m+\alpha,\frac{2+m+\alpha}{2}}([\varepsilon,1] \times [\varepsilon,T]), \quad \text{for each fixed } \varepsilon > 0.$$
 (1.62)

Therefore, by the definition of the spaces $H^{\lambda,\frac{\lambda}{2}}$ (see Appendix C), we have that

 $v_y \in H^{1+m+\alpha,\frac{1+m+\alpha}{2}}([\varepsilon,1] \times [\varepsilon,T]),$ for each fixed $\varepsilon > 0.$ (1.63) Thus, taking (1.58) into account,

$$\dot{s} \in H^{1+m+\alpha,\frac{1+m+\alpha}{2}}([\varepsilon,1]\times[\varepsilon,T]), \quad \text{for each fixed } \varepsilon > 0 \quad (1.64)$$

(we have used also (C.1) of Appendix C). The induction step, and the proof, are completed. $\hfill \Box$

CHAPTER 2

Weak formulation of the Stefan problem

In this chapter we consider the weak formulation of the Stefan problem. As in other PDE problems, the weak formulation actually takes the form of an integral equality. It is to be noted that any *explicit* reference to the free boundary is dropped from the weak formulation.

We'll comment on the modeling differences between the classical and the weak formulations, and give the basic mathematical results for the latter.

We work in the multi-dimensional case of a spatial domain $\Omega \subset \mathbb{R}^N$. We still denote the temperature by u in this chapter.

2.1. From the energy balance to the weak formulation

The heat equation

$$u_t = \operatorname{div}(\nabla u) + f, \qquad (2.1)$$

amounts to an energy balance equating the local change in time of 'energy' (expressed by u_t ; various physical constants are normalized to 1 here), to the divergence of the 'energy flux', plus the contribution of volumetric sources, represented by f.

The weak formulation is based on the extension of this idea to the case where the energy exhibits a jump at the critical temperature, due to the change of phase, as shown in Figure 2.1. Then we write (formally)

$$\frac{\partial}{\partial t}v = \operatorname{div}(\nabla u) + f, \qquad (2.2)$$

where the 'energy' (more exactly, the enthalpy) v jumps at the change of



FIGURE 2.1. Enthalpy v as a graph of temperature u, and viceversa.

phase. Specifically, solid at the critical temperature u = 0 corresponds to v = 0, while liquid at temperature u = 0 corresponds to v = 1 (we assume

the latent heat is normalized to unity). Where 0 < v < 1, therefore, change of phase is taking place, and the corresponding region is filled with a material whose state is neither pure solid nor pure liquid. Such regions are usually called *mushy regions*.

The standard heat equation is assumed to hold in the pure phases (i.e., where v > 1 or v < 0). This essentially amounts to

$$v = \begin{cases} u, & u < 0, \\ u+1, & u > 0. \end{cases}$$
(2.3)

Where 0 < v < 1, u must equal the critical temperature u = 0. It is therefore convenient to express the relation between v and u as follows

$$u(x,t) = \begin{cases} v(x,t), & v(x,t) \le 0, \\ 0, & 0 < v(x,t) < 1, \\ v(x,t) - 1, & v(x,t) \ge 1. \end{cases}$$
(2.4)

Note that v (not u) carries all the information on the state of the material. We can rephrase (2.4) in the language of graphs:

$$v \in E(u), \tag{2.5}$$

where E is the graph defined by

$$E(s) = \begin{cases} s, & s < 0, \\ [0,1], & s = 0, \\ s+1, & s > 0. \end{cases}$$
(2.6)

When v, u satisfy (2.5), we say that v is an admissible enthalpy for u, or that u is an admissible temperature for v.

Obviously, (2.2) can not be given a classical pointwise interpretation, since v is general not continuous (see (2.5)). Following an usual procedure, we obtain the *weak formulation* of (2.2) on multiplying both sides of it by a *testing function* $\varphi \in C_0^{\infty}(Q_T)$, and integrating (formally) by parts. In this way some of the derivatives appearing in (2.2) are unloaded on the smooth testing function. We obtain

$$\iint_{Q_T} \{-v\varphi_t + \nabla u \cdot \nabla \varphi\} \,\mathrm{d}x \,\mathrm{d}t = \iint_{Q_T} f\varphi \,\mathrm{d}x \,\mathrm{d}t \,. \tag{2.7}$$

Note that this formulation requires only we give a meaning to the first spatial derivatives of u (for example, u may be a Sobolev function). The complete formulation of the Stefan problem will be given below (see Section 2.2). The notion of weak solutions to the Stefan problem was introduced in [14], [9].

2.2. Definition of weak solution

Let us define the 'inverse' of the graph E in (2.6). This is the function ϑ given by

$$\vartheta(r) = \begin{cases} r, & r \le 0, \\ 0, & 0 < r < 1, \\ r - 1, & 1 \le r. \end{cases}$$
(2.8)

Let $Q_T = \Omega \times (0, T)$, where $\Omega \subset \mathbf{R}^N$ is a bounded open set with boundary of class C^{∞} .

Let us consider the Stefan problem

$$v_t - \Delta \vartheta(v) = f(v), \quad \text{in } Q_T,$$
(2.9)

$$v(x,0) = v_0(x), \qquad x \in \Omega,$$
 (2.10)

$$\frac{\partial \vartheta(v)}{\partial \nu} = 0, \qquad \text{on } \partial \Omega \times (0, T), \qquad (2.11)$$

where $v_0 \in L^{\infty}(\Omega)$ and $f \in L^{\infty}(\mathbf{R}) \cap C^{\infty}(\mathbf{R})$ are given functions. We assume that for a fixed $\mu > 0$

$$|f(v_1) - f(v_2)| \le \mu |v_1 - v_2|, \quad \text{for all } v_1, v_2 \in \mathbf{R}.$$
 (2.12)

Definition 2.1. A function $v \in L^{\infty}(Q_T)$ is a weak solution to (2.9)–(2.11) if

$$u := \vartheta(v) \in L^2(0, T; W_1^2(\Omega)), \qquad (2.13)$$

and for all $\varphi \in W_1^2(Q_T)$ such that $\varphi(x,T) = 0$ we have

$$\iint_{Q_T} \{-v\varphi_t + \nabla u \cdot \nabla \varphi\} \, \mathrm{d}x \, \mathrm{d}t = \int_{\Omega} v_0(x)\varphi(x,0) \, \mathrm{d}x + \iint_{Q_T} f(v)\varphi \, \mathrm{d}x \, \mathrm{d}t \,. \tag{2.14}$$

Note that equation (2.14) is obtained integrating formally by parts equation (2.9), after multiplying it by a φ as above.

Of course the structure of the problem could be generalized; for example the smoothness required of $\partial \Omega$ can be reduced by approximating Ω with more regular domains (but (2.12) is going to play an essential role). However we aim here at showing some basic techniques in the simple setting above, which is suitable for our purposes.

Remark 2.2. Actually, if v is a weak solution to the Stefan problem (in a sense similar to ours), the corresponding temperature u is continuous in Q_T (see [4]), a fact which however we won't use here.

2.2.1. Exercises.

Exercise 2.3. Assume $u \in C(Q_T)$ (see also Remark 2.2). Show that u is of class C^{∞} where it is not zero.

2.3. Comparing the weak and the classical formulations

2.3.1. The spatial normal. Let S be a smooth surface of \mathbf{R}^{N+1} , which we may assume for our purposes to be locally represented in the form

$$\Phi(x,t) = 0, \qquad (2.15)$$

with $\Phi \in C^1(\mathbb{R}^{N+1})$, and $\nabla \Phi \neq 0$ everywhere. Here we denote by $\nabla \Phi$ the gradient of Φ with respect to x, and by $\tilde{\nabla} \Phi = (\nabla \Phi, \Phi_t)$ the complete gradient of Φ with respect to (x, t).

We may think of S as of a moving surface in \mathbb{R}^{N} . More exactly, at each fixed instant t the surface takes the position

$$S(t) = \{x \in \mathbf{R}^N \mid \Phi(x, t) = 0\}.$$

A moving point x(t) belongs to S(t) for all t if and only if

$$\Phi(x(t), t) = 0$$
, for all t ,

which is equivalent, up to the choice of suitable initial data, to

$$\nabla \varPhi(x(t),t) \cdot \dot{x}(t) + \varPhi_t(x(t),t) = 0, \qquad \text{for all } t.$$

Define the spatial normal on S by

$$\nu = \frac{\nabla \Phi(x(t), t)}{|\nabla \Phi(x(t), t)|}.$$
(2.16)

The spatial normal, of course, is defined up to a change in sign. We have for all motions $t \mapsto x(t)$ as above

$$\dot{x}(t) \cdot \nu = -\frac{\Phi_t(x(t), t)}{|\nabla \Phi(x(t), t)|}.$$

This shows that, at a given position on S, the component of the velocity \dot{x} along the spatial normal is independent of the motion x. This quantity is referred to as the normal velocity V of S. Therefore, we have by definition

$$V(x,t) = -\frac{\Phi_t(x(t),t)}{|\nabla \Phi(x(t),t)|}$$

Again, note that V is defined up to a change in sign.

The complete normal to S at (x, t) is clearly $\boldsymbol{\nu} = (\boldsymbol{\nu}_x, \nu_t)$, where

$$\begin{split} \boldsymbol{\nu}_x &= \frac{\nabla \varPhi(x,t)}{|\tilde{\nabla}\varPhi(x,t)|} = \nu \frac{|\nabla \varPhi(x,t)|}{|\tilde{\nabla}\varPhi(x,t)|} \,,\\ \boldsymbol{\nu}_t &= \frac{\varPhi_t(x,t)}{|\tilde{\nabla}\varPhi(x,t)|} \,. \end{split}$$

2.3.2. Smooth weak solutions, with smooth interfaces, are classical solutions. Let us assume that a function u satisfies

$$u > 0, \qquad \text{in } A, \\ u < 0, \qquad \text{in } B,$$

and that u = 0 on the common portion of the boundaries of the open sets A, $B \subset \mathbb{R}^{N+1}$. We assume this portion to be a smooth surface S, with complete normal $\boldsymbol{\nu} = (\boldsymbol{\nu}_x, \nu_t)$ and spatial normal $\boldsymbol{\nu}$, according to the notation above. Let $\boldsymbol{\nu}$ be the outer normal to B. Moreover assume

$$u, f \in C(\overline{A \cup B}), \qquad u_{|A} \in C^{2,1}(\overline{A}), \qquad u_{|B} \in C^{2,1}(\overline{B}).$$

Finally, assume v (defined as in (2.3)) is a weak solution of (2.7), for any smooth φ whose support is contained in the interior of $A \cup B \cup S$. It will be apparent from our calculations that the definition of v at u = 0 is not relevant, in this case, essentially because the N + 1-dimensional measure of the free boundary u = 0 is zero.

By direct calculation we have, owing to the regularity of u and of v,

$$\iint_{A} v\varphi_t \, \mathrm{d}x \, \mathrm{d}t = -\int_{S} \varphi E(0+)\nu_t \, \mathrm{d}\sigma - \iint_{A} v_t \varphi \, \mathrm{d}x \, \mathrm{d}t$$
$$\iint_{B} v\varphi_t \, \mathrm{d}x \, \mathrm{d}t = \int_{S} \varphi E(0-)\nu_t \, \mathrm{d}\sigma - \iint_{B} v_t \varphi \, \mathrm{d}x \, \mathrm{d}t \, .$$

On adding these two equalities we find, recalling the definition of E,

$$\iint_{A\cup B} v\varphi_t \,\mathrm{d}x \,\mathrm{d}t = -\int_{S} \varphi \nu_t \,\mathrm{d}\sigma - \iint_{A\cup B} u_t \varphi \,\mathrm{d}x \,\mathrm{d}t \,, \tag{2.17}$$

since $v_t = u_t$ both in A and in B. The space part of the differential operator in (2.7) is treated similarly

$$\iint_{A} \nabla u \cdot \nabla \varphi \, \mathrm{d}x \, \mathrm{d}t = -\int_{S} \nabla u \cdot \boldsymbol{\nu}_{x} \, \mathrm{d}\sigma - \iint_{A} \varphi \, \Delta u \, \mathrm{d}x \, \mathrm{d}t \, \mathrm{$$

Again, on adding these two equalities we find

$$\iint_{A\cup B} \nabla u \cdot \nabla \varphi \, \mathrm{d}x \, \mathrm{d}t = \int_{S} [\nabla u^B - \nabla u^A] \cdot \boldsymbol{\nu}_x \, \mathrm{d}\sigma - \iint_{A\cup B} \varphi \, \Delta u \, \mathrm{d}x \, \mathrm{d}t \,, \quad (2.18)$$

where we denote by $\nabla u^A \ [\nabla u^B]$ the trace on S of the spatial gradient of the restriction of u to A [B]. Combining (2.17) with (2.18) we arrive at

$$\iint_{A\cup B} f\varphi \,\mathrm{d}x \,\mathrm{d}t = \iint_{A\cup B} \{-v\varphi_t + \nabla u \cdot \nabla \varphi\} \,\mathrm{d}x \,\mathrm{d}t = \int_{S} \varphi[\nu_t + \nabla u^B \cdot \boldsymbol{\nu}_x - \nabla u^A \cdot \boldsymbol{\nu}_x] \,\mathrm{d}\sigma$$
$$+ \iint_{A\cup B} \varphi\{u_t - \Delta u\} \,\mathrm{d}x \,\mathrm{d}t \,. \quad (2.19)$$

Taking an arbitrary smooth φ supported in A, we immediately find that in A

$$u_t - \Delta u = f. \tag{2.20}$$

Of course the same PDE holds in B, by the same token. Hence, we may drop the last integral in (2.19). Then take $\varphi = \varphi_{\varepsilon}$, where for all $\varepsilon > 0$

$$\varphi_{\varepsilon|S} = \psi \in C_0^1(S); \quad |\varphi_{\varepsilon}| \le 1; \quad |\operatorname{supp} \varphi_{\varepsilon}|_{N+1} \to 0, \quad \text{as } \varepsilon \to 0.$$

Then, taking $\varepsilon \to 0$ in (2.19) we get

$$\int_{S} \psi[\nu_t + \nabla u^B \cdot \boldsymbol{\nu}_x - \nabla u^A \cdot \boldsymbol{\nu}_x] \, \mathrm{d}\sigma = 0.$$

As ψ is reasonably arbitrary, it follows that on S

$$\nabla u^B \cdot \boldsymbol{\nu}_x - \nabla u^A \cdot \boldsymbol{\nu}_x = -\nu_t \,.$$

This condition can be rewritten as

$$V = \nabla u^B \cdot \nu - \nabla u^A \cdot \nu, \qquad (2.21)$$

23

,

and is the multi-dimensional equivalent of the Stefan condition (1.5) of Chapter 1. In fact, it could be directly derived from an energetical balance argument, as we did for (1.5) of Chapter 1. In this last approach, the weak formulation of the Stefan problem follows from (2.21) and from the heat equation which we assume to hold in A and in B separately: we only need go over our previous calculations in reverse order.

The case where either $v \equiv 1$ in A or $v \equiv 0$ in B can be treated similarly.

2.3.3. Some smooth weak solutions are not classical solutions. Let us consider the following problem

$$v_t - u_{xx} = 1$$
, in $Q_T = (0, 1) \times (0, +\infty)$, (2.22)

$$u(x,0) = -1, \qquad 0 < x < 1,$$
(2.23)

$$u_x(0,t) = 0, \qquad 0 < t,$$
 (2.24)

$$u_x(1,t) = 0, \qquad 0 < t$$
 (2.25)

(see [15]). We perform only a local analysi of the problem in the interior of the domain Q_T , giving for granted the solutions below actually take the boundary data (in a suitable sense).

If, instead of (2.22), the standard heat equation

$$u_t - u_{xx} = 1$$

was prescribed, clearly the solution to the initial value boundary problem would be

$$u(x,t) = -1 + t$$
, $0 < x < 1$, $0 < t$.

Let us check that this function can not be a solution to the Stefan problem above. Otherwise, we would have

$$v(x,t) = -1 + t \,, \quad 0 < t < 1 \,; \qquad v(x,t) = t \,, \quad 1 < t \,.$$

Thus for every $\varphi \in C_0^{\infty}(Q_T)$,

$$\iint_{Q_T} \{-v\varphi_t + u_x\varphi_x\} \,\mathrm{d}x \,\mathrm{d}t = -\iint_{Q_T} v\varphi_t \,\mathrm{d}x \,\mathrm{d}t = \iint_{Q_T} \varphi \,\mathrm{d}x \,\mathrm{d}t + \int_0^1 \varphi(x,1) \,\mathrm{d}x \,.$$

The last integral in this equality is evidently spurious, on comparison with the weak formulation (2.7).

Let us instead check that a solution (actually the unique solution, see Section 2.4) to (2.22)-(2.25) is given by

$$v(x,t) = -1 + t, \quad 0 < t; \qquad u(x,t) = \begin{cases} -1 + t, & 0 < t < 1, \\ 0, & 1 < t < 2, \\ -2 + t, & 2 < t. \end{cases}$$

It is immediately checked that (2.7) is fulfilled. We still have to check that (2.5) holds, that is that v is an admissible enthalpy for u. Again, this follows immediately from the definitions.

The above can be interpreted as follows: the enthalpy v grows in time accordingly to the prescribed volumetric source; the change of phase takes place over the time interval 1 < t < 2, because this is the time interval where $v \in (0, 1)$; over this time interval, therefore, the temperature equals

the critical temperature u = 0; for all other times, (2.22) coincides with the standard heat equation, and thus u is simply the solution to a suitable problem for the heat equation.

Note that the set u = 0 has in this example positive measure, in contrast with the classical formulation of Chapter 1. See also [8], [1] for a discussion of existence and non existence of mushy regions in weak solutions to change of phase problems.

2.3.4. Without sign restrictions, classical solutions may not be weak solutions. Let us go back to problem (1.16)-(1.21) of Chapter 1, where we now assume $u_0 \in C^1([0,b])$, $u_0(b) = 0$, $u_0(x) < 0$ for $0 \le x < b$, and, e.g., $h \equiv 0$. It can be shown (see [6]) that this problem has a classical solution, in the sense of Definition 1.8 of Chapter 1, at least for a small enough T > 0. Note that the Stefan condition (1.19) of Chapter 1 has now the 'wrong' sign (cf. Exercise 1.6 of Chapter 1). Therefore the classical solution at hand is *not* a solution of the weak formulation. Indeed, otherwise it would be a smooth weak solution with a smooth free boundary, and we would be able to infer the Stefan condition as above. However, as shown above, this condition would be the one 'correctly' corresponding to the actual sign of the solution, and therefore would be *different* from the one we prescribed.

More generally, no undercooling is possible in the weak formulation introduced here. In fact, the liquid and the solid phases are identified solely by the value of v. Thus, whenever u changes its sign a change of phase must take place. This is not the case in the classical formulation, where the liquid and solid phases are essentially identified by a topological argument, as the two connected components of the domain, separated by the special level surface which is *defined* as the free boundary. Other level surfaces corresponding to the value u = 0 may exist inside both phases.

2.3.5. Exercises.

Exercise 2.4. Assume that the interior M of the region $\{u = 0\}$ is non empty, where u is given by (2.4), and v satisfies (2.7). Show that, in a suitable weak sense, $v_t = f$ in M.

2.4. Uniqueness of the weak solution

2.4.1. A different notion of weak solution. If we assume in (2.14) that

i)
$$\varphi \in W_{2,1}^2(Q_T)$$
; ii) $\varphi(x,T) = 0$; iii) $\frac{\partial \varphi}{\partial \nu} = 0$, on $\partial \Omega \times (0,T)$;
(2.26)

we immediately obtain, on integrating once by parts,

$$-\iint_{Q_T} \{v\varphi_t + u\,\Delta\,\varphi\}\,\mathrm{d}x\,\mathrm{d}t = \int_{\Omega} v_0(x)\varphi(x,0)\,\mathrm{d}x + \iint_{Q_T} f(v)\varphi\,\mathrm{d}x\,\mathrm{d}t\,. \quad (2.27)$$

We need drop the requirement $\varphi(x,T) = 0$, for technical reasons. This can be done as follows. Choose a φ satisfying i), iii) of (2.26), but not necessarily ii). Fix $t \in (0, T)$, and define for $0 < \varepsilon < t$ (see Figure 2.2)

$$\chi_{\varepsilon}(\tau) = \min\left(1, \frac{1}{\varepsilon}(t-\tau)_+\right).$$

The function $\varphi \chi_{\varepsilon}$ satisfies requirement (2.26) in full, so that it can be taken



FIGURE 2.2. The auxiliary function χ_{ε} .

as a testing function in (2.27). Let us rewrite it as

$$-\iint_{Q_T} v[\varphi\chi_{\varepsilon}]_{\tau} \,\mathrm{d}x \,\mathrm{d}\tau = \int_{\Omega} v_0(x)\varphi(x,0) \,\mathrm{d}x + \iint_{Q_T} \{u\,\Delta\,\varphi + f(v)\varphi\}\chi_{\varepsilon} \,\mathrm{d}x \,\mathrm{d}\tau\,.$$

The behaviour of the right hand side as $\varepsilon \to 0$ is obvious. The left hand side equals

$$-\iint_{Q_T} v\varphi_\tau \chi_\varepsilon \,\mathrm{d}x \,\mathrm{d}\tau + \frac{1}{\varepsilon} \int_{t-\varepsilon}^t \int_{\Omega} v(x,\tau)\varphi(x,\tau) \,\mathrm{d}x \,\mathrm{d}\tau \,.$$

On letting $\varepsilon \to 0$ in this quantity we get, for almost all $t \in (0,T)$

$$-\iint_{Q_t} v\varphi_\tau \,\mathrm{d}x \,\mathrm{d}\tau + \int_{\Omega} v(x,t)\varphi(x,t) \,\mathrm{d}x \,.$$

Thus the definition of weak solution given above actually implies the new (and weaker) one

Definition 2.5. A function $v \in L^{\infty}(Q_T)$ is a weak solution of class L^{∞} to (2.9)–(2.11) if for all $\varphi \in W^2_{2,1}(Q_T)$ such that $\frac{\partial \varphi}{\partial \nu} = 0$ on $\partial \Omega \times (0,T)$ we have

$$\int_{\Omega} v(x,t)\varphi(x,t) \,\mathrm{d}x - \iint_{Q_t} \{ v\varphi_\tau + u\,\Delta\,\varphi \} \,\mathrm{d}x \,\mathrm{d}\tau$$
$$= \int_{\Omega} v_0(x)\varphi(x,0) \,\mathrm{d}x + \iint_{Q_t} f(v)\varphi \,\mathrm{d}x \,\mathrm{d}\tau \,, \quad (2.28)$$

for almost all $t \in (0, T)$. Here $u = \vartheta(v)$.

26

Note that we dropped in Definition 2.5 any regularity requirement for v (excepting boundedness).

Remark 2.6. It follows from (2.28) that the function

$$t\mapsto \int\limits_{\varOmega} v(x,t)\varphi(x,t)\,\mathrm{d}x$$

is actually continuous over [0, T], up to modification of v over sets of zero measure.

2.4.2. Continuous dependence on the initial data. We are now in a position to prove

Theorem 2.7. (CONTINUOUS DEPENDENCE ON THE DATA) Let v_1 , v_2 be two weak solutions of class L^{∞} to (2.9)–(2.11), in the sense of Definition 2.5 (or two weak solutions in the sense of Definition 2.1), corresponding to bounded initial data v_{01} , v_{02} respectively. Then, for almost all 0 < t < T

$$\int_{\Omega} |v_1(x,t) - v_2(x,t)| \, \mathrm{d}x \le e^{\mu t} \int_{\Omega} |v_{01}(x) - v_{02}(x)| \, \mathrm{d}x \,. \tag{2.29}$$

Corollary 2.8. (UNIQUENESS) Let v_1 , v_2 be two weak solutions of class L^{∞} to (2.9)–(2.11), in the sense of Definition 2.5 (or two weak solutions in the sense of Definition 2.1), corresponding to the same bounded initial data. Then $v_1 \equiv v_2$ in Q_T .

Remark 2.9. (SOLUTIONS OF CLASS L^1) Both Theorem 2.7 and its immediate Corollary 2.8 actually hold for a more general class of weak solutions, obtained replacing the requirement $v \in L^{\infty}(Q_T)$ in Definition 2.5 with $v \in L^1(Q_T)$. Also the initial data may be selected out of $L^1(\Omega)$. In this connection, in order to keep the integrals in (2.28) meaningful, we have to assume that φ is a Lipschitz continuous function in Q_T , with $\varphi_{x_i x_j} \in L^{\infty}(Q_T)$, $i, j = 1, \ldots, N$, and $\frac{\partial \varphi}{\partial \nu} = 0$ on $\partial \Omega \times (0, T)$.

Once existence of solutions in the sense of Definition 2.5 has been obtained, existence of solutions of class L^1 can be proven as follows. Assume $v_0 \in L^1(\Omega)$, and $v_0^i \to v_0$ in $L^1(\Omega)$, $v_0^i \in L^{\infty}(\Omega)$. Note that the solutions v^i of class L^{∞} corresponding to the approximating initial data v_0^i satisfy (2.29). Therefore $\{v^i\}$ is a Cauchy sequence in $L^1(Q_T)$, and we may assume it converges to a $v \in L^1(Q_T)$ both in the sense of $L^1(Q_T)$, and a.e. in Q_T . It is now a trivial task to take the limit in the weak formulation (2.28) satisfied by v^i and obtain the corresponding formulation for v.

Standard references for the material in this Section are [11], Chapter V, Section 9, and [12], whose approach we follow, with some modifications.

2.4.3. Proof of Theorem 2.7. Fix $t \in (0,T)$. Subtract from each other the two equations (2.28) written for the two solutions, and obtain

$$\int_{\Omega} [v_1(x,t) - v_2(x,t)]\varphi(x,t) \,\mathrm{d}x - \iint_{Q_t} (v_1 - v_2)[\varphi_\tau + a(x,t)\,\Delta\,\varphi] \,\mathrm{d}x \,\mathrm{d}\tau$$
$$= \int_{\Omega} [v_{01}(x) - v_{02}(x)]\varphi(x,0) \,\mathrm{d}x + \iint_{Q_t} [f(v_1) - f(v_2)]\varphi \,\mathrm{d}x \,\mathrm{d}\tau, \quad (2.30)$$

where we set

$$\begin{aligned} a(x,t) &= \frac{\vartheta(v_1(x,t)) - \vartheta(v_2(x,t))}{v_1(x,t) - v_2(x,t)} \,, \qquad v_1(x,t) \neq v_2(x,t) \,, \\ a(x,t) &= 0 \,, \qquad \qquad v_1(x,t) = v_2(x,t) \,. \end{aligned}$$

Due to the definition of ϑ we have

$$0 \le a(x,t) \le 1$$
, in Q_T . (2.31)

Next choose $\varphi = \varphi_{\varepsilon}$, where for each $\varepsilon > 0$, φ_{ε} is the solution of

$$\varphi_{\varepsilon\tau} + (a_{\varepsilon}(x,t) + \varepsilon) \Delta \varphi_{\varepsilon} = 0, \qquad \text{in } Q_t, \qquad (2.32)$$

$$\varphi(x,t) = \Phi(x), \qquad x \in \Omega,$$
 (2.33)

$$\frac{\partial \varphi}{\partial \nu} = 0,$$
 on $\partial \Omega \times (0, t).$ (2.34)

Here $\Phi \in C_0^{\infty}(\Omega), |\Phi(x)| \leq 1$, and $a_{\varepsilon} \in C^{\infty}(\overline{Q_T})$ satisfies

$$0 \le a_{\varepsilon} \le 1$$
, a.e. in Q_T ; $||a_{\varepsilon} - a||_2 \le \varepsilon$. (2.35)

Some relevant properties of φ_{ε} are collected in Lemma 2.10 below. Note that (2.33) is the initial value for the 'reverse' parabolic problem solved by φ_{ε} . By virtue of (2.40) we have as $\varepsilon \to 0$

$$\iint_{Q_t} \varepsilon |\Delta \varphi_{\varepsilon}| \, \mathrm{d}x \, \mathrm{d}\tau \leq \left(\iint_{Q_t} \varepsilon \, \mathrm{d}x \, \mathrm{d}\tau \right)^{1/2} \left(\iint_{Q_t} \varepsilon (\Delta \varphi_{\varepsilon})^2 \, \mathrm{d}x \, \mathrm{d}\tau \right)^{1/2} \\ \leq \sqrt{2t\varepsilon} \|\nabla \Phi\|_{2,\Omega} |\Omega|^{1/2} \to 0 \,,$$

as well as (using (2.35))

$$\begin{split} \iint_{Q_t} |a_{\varepsilon} - a| |\Delta \varphi_{\varepsilon}| \, \mathrm{d}x \, \mathrm{d}\tau &\leq \bigg(\iint_{Q_t} \frac{|a_{\varepsilon} - a|^2}{\varepsilon} \, \mathrm{d}x \, \mathrm{d}\tau \bigg)^{1/2} \bigg(\iint_{Q_t} \varepsilon (\Delta \varphi_{\varepsilon})^2 \, \mathrm{d}x \, \mathrm{d}\tau \bigg)^{1/2} \\ &\leq \sqrt{2\varepsilon} \|\nabla \Phi\|_{2,\Omega} \to 0 \,. \end{split}$$

Moreover, using (2.32) in (2.30), and (2.41), we get

$$\begin{split} \int_{\Omega(t)} & [v_1 - v_2] \varPhi \, \mathrm{d}x = \iint_{Q_t} [a - a_\varepsilon - \varepsilon] \, \Delta \, \varphi_\varepsilon [v_1 - v_2] \, \mathrm{d}x \, \mathrm{d}\tau + \int_{\Omega} [v_{01} - v_{02}] \varphi_\varepsilon (x, 0) \, \mathrm{d}x \\ & + \iint_{Q_t} [f(v_1) - f(v_2)] \varphi_\varepsilon \, \mathrm{d}x \, \mathrm{d}\tau \leq (\|v_1\|_\infty + \|v_2\|_\infty) \iint_{Q_t} [|a - a_\varepsilon| + \varepsilon] |\Delta \, \varphi_\varepsilon | \, \mathrm{d}x \, \mathrm{d}\tau \\ & + \int_{\Omega} |v_{01} - v_{02}| \, \mathrm{d}x + \mu \iint_{Q_t} |v_1 - v_2| \, \mathrm{d}x \, \mathrm{d}\tau \, . \end{split}$$

As $\varepsilon \to 0$ this yields

$$\int_{\Omega(t)} [v_1 - v_2] \Phi \, \mathrm{d}x \le \int_{\Omega} |v_{01} - v_{02}| \, \mathrm{d}x + \mu \int_{0}^{t} \int_{\Omega(\tau)} |v_1 - v_2| \, \mathrm{d}x \, \mathrm{d}\tau \,.$$
(2.36)

Choose now $\Phi = \Phi_n$, where for $n \to \infty$

$$\Phi_n(x) \to \operatorname{sign}\left(v_1(x,t) - v_2(x,t)\right), \quad \text{a.e. } x \in \Omega.$$

On letting $n \to \infty$ in (2.36) we obtain

$$\int_{\Omega(t)} |v_1 - v_2| \, \mathrm{d}x \le \int_{\Omega} |v_{01} - v_{02}| \, \mathrm{d}x + \mu \int_{0}^{t} \int_{\Omega(\tau)} |v_1 - v_2| \, \mathrm{d}x \, \mathrm{d}\tau \, .$$

The statement now follows simply invoking Gronwall's lemma.

Lemma 2.10. Let $\alpha \in C^{\infty}(\overline{Q_T})$, $0 < \varepsilon \leq \alpha \leq \alpha_0$, where ε and α_0 are given constants. Let $\Phi \in C_0^{\infty}(\Omega)$. Then there exists a unique solution $\varphi \in C^{\infty}(\overline{Q_T})$ of

$$\varphi_t - \alpha \Delta \varphi = 0, \qquad in \ Q_T, \qquad (2.37)$$

$$\varphi(x,0) = \Phi(x), \qquad x \in \Omega, \qquad (2.38)$$

$$\frac{\partial \varphi}{\partial \nu} = 0, \qquad on \ \partial \Omega \times (0, T), \qquad (2.39)$$

such that for all 0 < t < T

$$\iint_{Q_t} (\varphi_\tau^2 + \alpha (\Delta \varphi)^2) \, \mathrm{d}x \, \mathrm{d}\tau + \int_{\Omega(t)} |\nabla \varphi|^2 \, \mathrm{d}x \le (\alpha_0 + 1) \int_{\Omega} |\nabla \Phi|^2 \, \mathrm{d}x \,. \quad (2.40)$$

Moreover

$$\|\varphi\|_{\infty} \le \|\Phi\|_{\infty} \,. \tag{2.41}$$

PROOF. The existence of a unique solution $\varphi \in C^{\infty}(\overline{Q_T})$ to (2.37)–(2.39) is a classical result. Let us multiply (2.37) by $\Delta \varphi$, and integrate by parts over Q_t , for an arbitrarily fixed $t \in (0, T)$. We find

$$\iint_{Q_t} \alpha |\Delta \varphi|^2 \, \mathrm{d}x \, \mathrm{d}\tau = \iint_{Q_t} \varphi_\tau \, \Delta \varphi \, \mathrm{d}x \, \mathrm{d}\tau = -\iint_{Q_t} \nabla \varphi_\tau \cdot \nabla \varphi \, \mathrm{d}x \, \mathrm{d}\tau$$
$$= \frac{1}{2} \iint_{\Omega} |\nabla \varphi(x,0)|^2 \, \mathrm{d}x - \frac{1}{2} \iint_{\Omega} |\nabla \varphi(x,t)|^2 \, \mathrm{d}x$$

Using again (2.37) we obtain

$$\iint_{Q_t} \varphi_{\tau}^2 \, \mathrm{d}x \, \mathrm{d}\tau = \iint_{Q_t} \alpha^2 |\Delta \varphi|^2 \, \mathrm{d}x \, \mathrm{d}\tau \le \frac{\alpha_0}{2} \int_{\Omega} |\nabla \varphi(x,0)|^2 \, \mathrm{d}x \, .$$

Thus, φ satisfies the integral estimate (2.40). The bound in (2.41) is an obvious consequence of the maximum and boundary point principles of Appendix A.

2.4.4. Exercises.

Exercise 2.11. Prove that

$$\Phi \ge 0 \ [\le 0] \Longrightarrow \varphi_{\varepsilon} \ge 0 \ [\le 0]$$

where φ_{ε} is the solution to (2.32)–(2.34). Use this fact to prove the comparison result

$$v_{01} \le v_{02}$$
 in $\Omega \Longrightarrow v_1 \le v_2$ in Q_T

where v_1 and v_2 are as in Theorem 2.7, provided $f' \ge 0$.

Exercise 2.12. If $\Phi \in C^{\infty}(\overline{\Omega})$, but $\Phi \notin C_0^{\infty}(\Omega)$, φ as in Lemma 2.10 need not be even $C^1(\overline{Q_T})$. Why?

2.5. Existence of weak solutions

We apply here the ideas of [8], though we approximate the Stefan problem with smooth parabolic problems, rather than discretizing it in time.

Theorem 2.13. There exists a weak solution v to (2.9)-(2.11), in the sense of Definition 2.1, satisfying

$$\|v\|_{\infty} \le \|v_0\|_{\infty} \,. \tag{2.42}$$

The proof of this existence result relies on an approximation procedure. Namely, we approximate (2.9)-(2.11) with a sequence of smoothed problems; the solutions to these problems in turn approach a solution to the original Stefan problem.

We need a sequence of smooth constitutive functions $\vartheta_n \in C^{\infty}(\mathbb{R}^N)$ approximating ϑ , such that

$$\frac{1}{n} \le \vartheta'_n(s) \le 1, \quad s \in \mathbf{R}; \qquad \vartheta_n \to \vartheta, \quad \text{uniformly in } \mathbf{R}.$$
(2.43)

Clearly we may assume

$$\vartheta(s) \le \vartheta_n(s) \le \vartheta(s) + \frac{1}{n}, \quad s \in \mathbf{R}; \qquad \vartheta(s) = \vartheta_n(s) = s, \quad s < 0.$$
 (2.44)

Define E_n as the inverse function of ϑ_n . Then

$$E_n(s) \le E(s), \quad s \in \mathbf{R}; \qquad E(s) = E_n(s) = s, \quad s < 0.$$
 (2.45)

Let us also introduce a sequence $v_{0n} \in C_0^{\infty}(\Omega)$ approximating the initial data as in

$$v_{0n} \to v_0$$
, a.e. in Ω ; $||v_{0n}||_{\infty} \le ||v_0||_{\infty}$. (2.46)


FIGURE 2.3. The approximating functions ϑ_n and E_n .

For each n there exists a unique solution $v_n \in C^{\infty}(\overline{Q_T})$ to

$$v_{nt} - \Delta \vartheta_n(v_n) = f(v_n)\eta_n, \quad \text{in } Q_T, \qquad (2.47)$$

$$\frac{\partial \vartheta_n(v_n)}{\partial \nu} = 0, \qquad \text{on } \partial \Omega \times (0,T), \qquad (2.48)$$

$$v_n(x,0) = v_{0n}(x), \quad \text{in } \Omega,$$
 (2.49)

where $\eta_n \in C_0^{\infty}(\Omega)$ is such that

$$\eta_n(x) = 1$$
, $\operatorname{dist}(x, \partial \Omega) > \frac{1}{n}$; $0 \le \eta_n(x) \le 1$, $x \in \Omega$.

Let us denote $u_n = \vartheta_n(v_n)$, and rewrite (2.47) as

$$E_n(u_n)_t - \Delta u_n = f(v_n)\eta_n, \quad \text{in } Q_T.$$
(2.50)

Owing to the maximum principle (see Theorem A.4 of Appendix A), we have

$$||u_n||_{\infty} \le ||v_n||_{\infty} \le ||v_0||_{\infty} + T||f||_{\infty} =: M.$$
(2.51)

2.5.1. The energy inequality. Multiply (2.50) by u_n and integrate by parts. Note that

$$E_n(u_n)_t u_n = \frac{\partial}{\partial t} \int_0^{u_n} E'_n(s) s \, \mathrm{d}s \,,$$

and that, if k > 0, taking into account (2.43),

$$\frac{k^2}{2} \le \int_0^k E'_n(s) s \, \mathrm{d}s \le E_n(k) k \le (k+1)k \,.$$

If k < 0 we simply have

$$\int_{0}^{k} E'_n(s)s \,\mathrm{d}s = \frac{k^2}{2} \,.$$

Therefore we obtain after standard calculations, for each $t \in (0, T)$

$$\begin{split} \frac{1}{2} \int_{\Omega} u_n(x,t)^2 \, \mathrm{d}x + \iint_{Q_t} |\nabla u_n|^2 \, \mathrm{d}x \, \mathrm{d}\tau \\ &\leq \frac{1}{2} \int_{\Omega} |\vartheta_n(v_{0n})| (|\vartheta_n(v_{0n})| + 1) \, \mathrm{d}x + \iint_{Q_t} f(v_n) \eta_n u_n \, \mathrm{d}x \, \mathrm{d}\tau \\ &\leq \frac{|\Omega|}{2} (\|v_0\|_{\infty} + 1)^2 + \|f\|_{\infty} M |\Omega| T \,. \end{split}$$

Note that both terms on the leftmost side of this estimate are positive. Dropping either one, taking the supremum in time, and collecting the two bounds so obtained, we get

$$\sup_{0 < t < T} \int_{\Omega} u_n(x,t)^2 \, \mathrm{d}x + \iint_{Q_T} |\nabla u_n|^2 \, \mathrm{d}x \, \mathrm{d}\tau \le C \,, \tag{2.52}$$

where C > 0 is a constant depending on the data of the problem, but not on n.

It follows that we may extract a subsequence, still labelled by n, such that

$$u_n \to u, \quad \nabla u_n \to \nabla u, \quad \text{weakly in } L^2(Q_T).$$
 (2.53)

Moreover we have

$$\|u\|_{\infty} \le M \,, \tag{2.54}$$

and

$$\sup_{0 < t < T} \int_{\Omega} u(x,t)^2 \, \mathrm{d}x + \iint_{Q_T} |\nabla u|^2 \, \mathrm{d}x \, \mathrm{d}\tau \le C \,. \tag{2.55}$$

We may as well assume

$$v_n \to v$$
, weakly in $L^2(Q_T)$, (2.56)

but note that, due to the nonlinear nature of our problem (i.e., the fact that E and f are not linear functions), weak convergence is not enough to pass to the limit in the weak formulation of the approximating problem, i.e., in

$$\iint_{Q_T} \{-v_n \varphi_t + \nabla u_n \cdot \nabla \varphi\} \,\mathrm{d}x \,\mathrm{d}t = \int_{\Omega} v_{0n}(x) \varphi(x,0) \,\mathrm{d}x + \iint_{Q_T} f(v_n) \eta_n \varphi \,\mathrm{d}x \,\mathrm{d}t \,,$$
(2.57)

where φ is any function out of $W_1^2(Q_T)$ with $\varphi(x,T) = 0$. For example we do not know that v is an admissible enthalpy for u. We must therefore obtain some stronger kind of convergence for the sequence v_n , so that, e.g., $f(v_n)$ converges to f(v).

However we do have some compactness in suitable integral norms for the sequence u_n , due to (2.52). More specifically, let $h \in \mathbb{R}^N$ be any given vector with length $0 < |h| < \delta$, and let $\mathbf{k} = h/|h|$. Setting

$$\Omega_{\delta} = \{ x \in \Omega \mid \operatorname{dist}(x, \partial \Omega) > \delta \},\$$

we calculate, by a standard argument,

$$\int_{0}^{T} \int_{\Omega_{\delta}} |u_{n}(x+h,t) - u_{n}(x,t)| \, \mathrm{d}x \, \mathrm{d}t = \int_{0}^{T} \int_{\Omega_{\delta}} \left| \int_{0}^{|h|} \nabla u_{n}(x+sk) \cdot k \, \mathrm{d}s \right| \, \mathrm{d}x \, \mathrm{d}t$$

$$\leq \int_{0}^{T} \int_{\Omega_{\delta}} \int_{0}^{|h|} |\nabla u_{n}(x+sk)| \, \mathrm{d}s \, \mathrm{d}x \, \mathrm{d}t = \int_{0}^{T} \int_{0}^{|h|} \int_{\Omega_{\delta}} |\nabla u_{n}(x+sk)| \, \mathrm{d}x \, \mathrm{d}s \, \mathrm{d}t$$

$$\leq \int_{0}^{T} \int_{0}^{|h|} \int_{\Omega} |\nabla u_{n}(x)| \, \mathrm{d}x \, \mathrm{d}s \, \mathrm{d}t = |h| \int_{0}^{T} \int_{\Omega} |\nabla u_{n}(x)| \, \mathrm{d}x \, \mathrm{d}t \leq C|h|. \quad (2.58)$$

We have used the fact that $x + s\mathbf{k} \in \Omega$ for all $x \in \Omega_{\delta}$ and all 0 < s < |h|, and (2.52).

2.5.2. The *BV* estimate. Introduce a cut off function $\zeta \in C^{\infty}(\Omega)$ such that

$$\begin{split} \zeta(x) &\equiv 1 \,, \quad \mathrm{dist}(x,\partial \Omega) > 4\delta \,; \qquad \zeta(x) \equiv 0 \,, \quad \mathrm{dist}(x,\partial \Omega) < 2\delta \,; \\ |\nabla \, \zeta| &\leq \frac{\gamma}{\delta} \,; \qquad |\Delta \, \zeta| \leq \frac{\gamma}{\delta^2} \,; \end{split}$$

where γ does not depend on δ . We may also assume that $n > 1/\delta$, so that $\eta_n \zeta \equiv \zeta$ in Ω .

In this Subsection we drop the index n, for ease of notation. Therefore we write ϑ for ϑ_n , v for v_n , and so on. For a given $h \in \mathbf{R}^N$ define the testing function

$$\varphi(x,t) = \operatorname{sign}_{\varepsilon} \left(\vartheta(v(x+h,t)) - \vartheta(v(x,t)) \right) \zeta(x)$$

where $\operatorname{sign}_{\varepsilon} \in C^{\infty}(\mathbf{R})$ is a smooth approximation of sign, such that

$$\operatorname{sign}_{\varepsilon}(s) \to \operatorname{sign}(s), \quad s \in \mathbf{R}, \qquad \operatorname{sign}'_{\varepsilon} \ge 0$$

(we set $\operatorname{sign}(0) = \operatorname{sign}_{\varepsilon}(0) = 0$). Let us denote

$$\varphi_h(x,t) = \varphi(x-h,t).$$

If $|h| < \delta$, which we assume from now on, both φ and φ_h vanish in a neighbourhood of $\partial \Omega \times (0,T)$.

Multiply (2.47) by φ and integrate by parts, obtaining

$$\iint_{Q_t} v_\tau \varphi \, \mathrm{d}x \, \mathrm{d}\tau + \iint_{Q_t} \nabla \vartheta(v) \cdot \nabla \varphi \, \mathrm{d}x \, \mathrm{d}\tau = \iint_{Q_t} f(v) \varphi \, \mathrm{d}x \, \mathrm{d}\tau \,. \tag{2.59}$$

On performing the same operation with φ_h , we obtain

$$\iint_{Q_t} v_{\tau}(x,\tau)\varphi(x-h,\tau) \,\mathrm{d}x \,\mathrm{d}\tau + \iint_{Q_t} \nabla \,\vartheta(v(x,\tau)) \cdot \nabla \,\varphi(x-h,\tau) \,\mathrm{d}x \,\mathrm{d}\tau$$
$$= \iint_{Q_t} f(v(x,\tau))\varphi(x-h,\tau) \,\mathrm{d}x \,\mathrm{d}\tau \,. \quad (2.60)$$

Let us change the integration variable in (2.60),

$$x - h \mapsto y$$
.

The domain of integration stays the same (i.e., Q_t) because

$$\operatorname{supp} \varphi_h \subset Q_T, \qquad \operatorname{supp} \varphi \subset Q_T.$$

Still denoting the new variable by x we obtain

$$\iint_{Q_t} v_{\tau}(x+h,\tau)\varphi(x,\tau) \,\mathrm{d}x \,\mathrm{d}\tau + \iint_{Q_t} \nabla \,\vartheta(v(x+h,\tau)) \cdot \nabla \,\varphi(x,\tau) \,\mathrm{d}x \,\mathrm{d}\tau$$
$$= \iint_{Q_t} f(v(x+h,\tau))\varphi(x,\tau) \,\mathrm{d}x \,\mathrm{d}\tau \,. \quad (2.61)$$

On subtracting (2.59) from (2.61), and explicitly calculating $\nabla\,\varphi,$ we arrive at

$$\iint_{Q_t} [v_{\tau}(x+h,\tau) - v_{\tau}(x,\tau)]_{\tau} \varphi(x,\tau) \, \mathrm{d}x \, \mathrm{d}\tau$$

$$+ \iint_{Q_t} |\nabla \vartheta(v(x+h,\tau)) - \nabla \vartheta(v(x,\tau))|^2 \operatorname{sign}_{\varepsilon}'(\dots) \zeta \, \mathrm{d}x \, \mathrm{d}\tau$$

$$+ \iint_{Q_t} [\nabla \vartheta(v(x+h,\tau)) - \nabla \vartheta(v(x,\tau))] \cdot \nabla \zeta \operatorname{sign}_{\varepsilon}(\dots) \, \mathrm{d}x \, \mathrm{d}\tau$$

$$= \iint_{Q_t} [f(v(x+h,\tau)) - f(v(x,\tau))] \varphi(x,\tau) \, \mathrm{d}x \, \mathrm{d}\tau . \quad (2.62)$$

The second integral in (2.62) may be dropped, since it is non negative. Next note

$$v(x+h,\tau) = v(x,\tau) \iff \vartheta(v(x+h,\tau)) = \vartheta(v(x,\tau)),$$

as $\vartheta=\vartheta_n$ is strictly increasing. Thus, as $\varepsilon\to 0$

$$\operatorname{sign}_{\varepsilon} \left(\vartheta(v(x+h,\tau)) - \vartheta(v(x,\tau)) \right) \to \operatorname{sign} \left(\vartheta(v(x+h,\tau)) - \vartheta(v(x,\tau)) \right) \\ = \operatorname{sign} \left(v(x+h,\tau) - v(x,\tau) \right) =: \sigma(x,\tau) \,.$$

Finally, we may take the limit $\varepsilon \to 0$ in (2.62), after dropping the second integral as we said, to find

$$\iint_{Q_t} [v_{\tau}(x+h,\tau) - v_{\tau}(x,\tau)]\sigma(x,\tau)\zeta(x) \,\mathrm{d}x \,\mathrm{d}\tau
+ \iint_{Q_t} [\nabla \vartheta(v(x+h,\tau)) - \nabla \vartheta(v(x,\tau))] \cdot \nabla \zeta \sigma(x,\tau) \,\mathrm{d}x \,\mathrm{d}\tau
\leq \iint_{Q_t} [f(v(x+h,\tau)) - f(v(x,\tau))]\sigma(x,\tau)\zeta \,\mathrm{d}x \,\mathrm{d}\tau. \quad (2.63)$$

The first integral in (2.63) equals

$$\iint_{Q_t} \frac{\partial}{\partial \tau} |v_{\tau}(x+h,\tau) - v_{\tau}(x,\tau)|\zeta(x) \,\mathrm{d}x \,\mathrm{d}\tau$$
$$= \int_{\Omega} |v(x+h,t) - v(x,t)|\zeta(x) \,\mathrm{d}x - \int_{\Omega} |v_0(x+h) - v_0(x)|\zeta(x) \,\mathrm{d}x \,.$$

The second integral in (2.63) equals

$$\begin{split} \iint_{Q_t} \nabla |\vartheta(v(x+h,\tau)) - \vartheta(v(x,\tau))| \cdot \nabla \zeta \, \mathrm{d}x \, \mathrm{d}\tau \\ &= - \iint_{Q_t} |\vartheta(v(x+h,\tau)) - \vartheta(v(x,\tau))| \, \Delta \zeta \, \mathrm{d}x \, \mathrm{d}\tau \,, \end{split}$$

so that, according to (2.58), its absolute value is majorised by

$$\frac{\gamma}{\delta^2} \iint_{\Omega_{\delta} \times (0,t)} |\vartheta(v(x+h,\tau)) - \vartheta(v(x,\tau))| \, \mathrm{d}x \, \mathrm{d}\tau \le \frac{C}{\delta^2} |h| \, .$$

The third and last integral in (2.63) is bounded simply by taking into account the Lipschitz continuity of f. Collecting these estimates we find

$$\int_{\Omega} |v(x+h,t) - v(x,t)|\zeta(x) \, \mathrm{d}x \le \int_{\Omega} |v_0(x+h) - v_0(x)|\zeta(x) \, \mathrm{d}x + \frac{C}{\delta^2} |h| + \mu \int_0^t \int_{\Omega} |v(x+h,\tau) - v(x,\tau)|\zeta(x) \, \mathrm{d}x \, \mathrm{d}\tau \,. \tag{2.64}$$

By Gronwall's lemma we conclude that

$$\int_{\Omega} |v(x+h,t) - v(x,t)|\zeta(x) \,\mathrm{d}x \le e^{\mu t} \left\{ \int_{\Omega} |v_0(x+h) - v_0(x)|\zeta(x) \,\mathrm{d}x + \frac{C}{\delta^2} |h| \right\} \quad (2.65)$$

Assume now that the original initial data v_0 satisfies for all $h \in \mathbf{R}^N$, $|h| < \delta$, $\delta > 0$,

$$\int_{\Omega_{\delta}} |v_0(x+h) - v_0(x)| \,\mathrm{d}x \le C_{\delta} |h| \,. \tag{2.66}$$

We may therefore assume the approximating initial data v_{0n} satisfy a similar inequality (see Lemma B.1 of Appendix B). This and (2.65) allow us to conclude that for all 0 < t < T, $h \in \mathbf{R}^N$, $n \ge 1$,

$$\int_{\Omega_{\delta}} |v_n(x+h,t) - v_n(x,t)| \, \mathrm{d}x \le C_{\delta} |h| \,. \tag{2.67}$$

Standard (and trivial) results in functional analysis imply that (2.67) is equivalent, since $v_n \in C^{\infty}(\overline{Q_T})$, to

$$\int_{\Omega_{\delta}} |\nabla v_n(x,t)| \, \mathrm{d}x \le C_{\delta} \,. \tag{2.68}$$

Here and above, C_{δ} denotes a constant depending on δ , but not on n.

Remark 2.14. In general, the L^1 estimate (2.67) does not imply for an integrable function v the existence of the gradient ∇v in the sense of Sobolev. This is a marked difference with the case of similar L^p estimates, with p > 1. Instead, estimates like (2.67) imply that v is a function of bounded variation, or BV function, whence the title of this subsection.

2.5.3. Compactness of v_n in L^1 . In order to obtain the desired compactness of the sequence $\{v_n\}$, in $L^1_{loc}(Q_T)$, we still have to complement (2.67) with a similar estimate involving translations in *time* rather than in *space*. This bound will be achieved as a consequence of a theorem by Kruzhkov, which we state and prove in Appendix B. Essentially, the result states that if we already know some regularity of the solution to a parabolic equation at each time level, we may infer some (lesser) regularity in the time variable. The regularity in space is in our case guaranteed by (2.67). The remarkable input of the theorem is that strong continuity in an integral norm is a consequence of a notion of weak continuity.

Let $g = g(x), g \in C_0^1(\Omega_{\delta})$. On multiplying (2.47) by g and integrating over Ω between t and t + s, 0 < t < t + s < T, standard calculations give

$$\int_{\Omega} g(x)[v_n(x,t+s) - v_n(x,t)] \, \mathrm{d}x = - \int_{t}^{t+s} \int_{\Omega} \nabla g \cdot \nabla \vartheta_n(v_n) \, \mathrm{d}x \, \mathrm{d}\tau + \int_{t}^{t+s} \int_{\Omega} f(v_n) \eta_n g \, \mathrm{d}x \, \mathrm{d}\tau \,. \quad (2.69)$$

Note that

$$\left| \int_{t}^{t+s} \int_{\Omega} \nabla g \cdot \nabla \vartheta_{n}(v_{n}) \, \mathrm{d}x \, \mathrm{d}\tau \right| \leq \int_{t}^{t+s} \int_{\Omega} \vartheta_{n}'(v_{n}) |\nabla g| |\nabla v_{n}| \, \mathrm{d}x \, \mathrm{d}\tau$$
$$\leq \|\vartheta_{n}'\|_{\infty} \|\nabla g\|_{\infty} s \sup_{t < \tau < t+s} \int_{\Omega_{\delta}} |\nabla v_{n}(x,\tau)| \, \mathrm{d}x \leq C_{\delta} \|\nabla g\|_{\infty} s \,,$$

owing to estimate (2.68). Clearly C_{δ} does not depend on n. Therefore

$$\left| \int_{\Omega} g(x) [v_n(x,t+s) - v_n(x,t)] \,\mathrm{d}x \right| \le C_{\delta} \big(\|g\|_{\infty} + \|\nabla g\|_{\infty} \big) s \,, \qquad (2.70)$$

for all $g \in C_0^1(\Omega_{\delta})$. Then, Kruzhkov's Theorem B.3 of Appendix B (see also Remark B.4 there) implies that

$$\int_{\Omega_{\delta}} |v_n(x,t+s) - v_n(x,t)| \, \mathrm{d}x \le C_{\delta}\sqrt{s} \,, \qquad 0 < s < \delta^2 \,, \tag{2.71}$$

for all 0 < t < t + s < T, and for all $\delta > 0$. Combining (2.71) with (2.67) we obtain that the sequence $\{v_n\}$ is pre-compact in $L^1(\Omega_{\delta} \times (0,T))$. By means of usual diagonal procedures, we may extract a subsequence (still labelled by n) such that

$$v_n \to v$$
, a.e. in Q_T . (2.72)

It follows

$$u_n = \vartheta_n(v_n) \to \vartheta(v) = u$$
, a.e. in Q_T , (2.73)

where clearly u must equal the weak limit in (2.53).

For any testing function φ as in Definition 2.1, we may therefore take the limit $n \to \infty$ in (2.57), and obtain (2.14). By construction, v and u satisfy the relevant regularity requirements. By the same token, $\vartheta(v) = u$, i.e., $v \in E(u)$.

2.5.4. Removing the extra assumption on v_0 . We have so far proven existence of solutions under the extra regularity assumption (2.66), which is certainly satisfied, for example, if the initial data are in $C^1(\overline{\Omega})$. To extend the proof to the case when v_0 is merely assumed to be a bounded function (so that (2.66) does not necessarily hold), consider first a sequence of smoothed initial data $v_0^j \in C^1(\overline{\Omega})$, such that

$$v_0^j \to v_0$$
, in $L^1(\Omega)$, $||v_0^j||_{\infty} \le ||v_0||_{\infty}$. (2.74)

Consider the sequence v^j of weak solutions, corresponding to these initial data, to the Stefan problem. These are therefore solutions in the sense of Definition 2.1. Note that, perhaps extracting a subsequence we may still assume that v^j , $u^j = \vartheta(v^j)$, ∇u^j converge weakly, because estimates (2.51), (2.52) are in force for them, uniformly on j.

Due to Theorem 2.7, and to (2.74), the sequence v^j actually is a Cauchy sequence in $L^1(Q_T)$. Again, perhaps extracting a subsequence we may assume that

$$v^j \to v$$
, a.e. in Q_T . (2.75)

Hence, we may take the limit in the integral equation satisfied by v^{j} , i.e.,

$$\iint_{Q_T} \{ -v^j \varphi_t + \nabla u^j \cdot \nabla \varphi \} \, \mathrm{d}x \, \mathrm{d}t = \int_{\Omega} v_0^j(x) \varphi(x,0) \, \mathrm{d}x + \iint_{Q_T} f(v^j) \varphi \, \mathrm{d}x \, \mathrm{d}t \,,$$

finally proving that v solves the original Stefan problem.

2.5.5. Exercises.

Exercise 2.15. Prove in detail that (2.55) follows from (2.52).

Exercise 2.16. A much simpler proof of existence of weak solutions is available when f does not depend on v (but, e.g., on u). In this case, indeed it is enough to prove strong convergence for the sequence u_n . To obtain the needed compactness estimate, which actually amounts to a bound of $||u_{nt}||_2$

uniform in n, multiply (2.50) by u_{nt} , and use the properties of E_n . Go over the details of this approach; e.g., prove that the weak limit v of v_n is an admissible enthalpy for the strong limit u of u_n .

Exercise 2.17. Assume that (2.66) is replaced by

$$\int_{\Omega_{\delta}} |v_0(x+h) - v_0(x)| \, \mathrm{d}x \le C_{\delta} \omega(|h|) \,,$$

for a non decreasing non negative continuous function $\omega : \mathbf{R} \to \mathbf{R}, \, \omega(0) = 0$, such that $\omega(s) \geq s, 0 < s < 1$. Prove that the solution v satisfies for $|h| < \delta < 1,$

$$\int_{\Omega_{\delta}} |v(x+h,t) - v(x,t)| \, \mathrm{d}x \le C_{\delta} \omega(|h|) \,, \qquad \text{a.e. } t \in (0,T).$$

2.6. A comparison result

Theorem 2.18. Let v_0^1 , $v_0^2 \in L^{\infty}(\Omega)$, and f^1 , $f^2 \in C^{\infty}(\mathbf{R})$ satisfy (2.12). Denote by v^1 , v^2 the corresponding weak solutions to the Stefan problem. If $v_0^1 \ge v_0^2$ in Ω , and $f^1 \ge f^2$ in \mathbf{R} , then $v^1 \ge v^2$ in Q_T .

PROOF. As we already know a result of uniqueness of the solution, we may prove the statement by approximation. By the same token, we may assume the initial data are smooth, e.g., in $C^1(\overline{\Omega})$. Let

$$\{v_n^i\}_{n=1}^\infty, \quad i=1,2$$

be the two sequences constructed in Section 2.5 as solutions to the problems (2.47)–(2.49), with v_{0n} [f] replaced with v_{0n}^i [fⁱ] respectively. Subtract from each other the PDE solved by v_n^1 and v_n^2 , and multiply the

resulting equation by

$$\varphi_{\varepsilon} = \operatorname{sign}_{\varepsilon}^{+}(\vartheta_n(v_n^2) - \vartheta_n(v_n^1))$$

where $\operatorname{sign}_{\varepsilon}^{+}$ is a smooth approximation of the function $\operatorname{sign}^{+}(s) = \chi_{(0,+\infty)}(s)$, with $\operatorname{sign}_{\varepsilon}^{+\prime} \geq 0$. Integrating by parts over Q_t we obtain

$$\iint_{Q_t} \left[(v_{n\tau}^2 - v_{n\tau}^1) \varphi_{\varepsilon} + \operatorname{sign}_{\varepsilon}^{+\prime} (u_n^2 - u_n^1) |\nabla (u_n^2 - u_n^1)|^2 \right] \mathrm{d}x \,\mathrm{d}\tau$$
$$= \iint_{Q_t} \left[f^2 (v_n^2) - f^1 (v_n^1) \right] \eta_n \varphi_{\varepsilon} \,\mathrm{d}x \,\mathrm{d}\tau.$$

On dropping the quadratic term above, and letting $\varepsilon \to 0$ we get (recall that $\operatorname{sign}^+(u_n^2 - u_n^1) = \operatorname{sign}^+(v_n^2 - v_n^1)$, see the discussion in Subsection 2.5.2)

$$\int_{\Omega} (v_n^2 - v_n^1)_+(x, t) \, \mathrm{d}x \le \iint_{Q_t} \left[f^2(v_n^2) - f^1(v_n^1) \right] \eta_n \operatorname{sign}^+(v_n^2 - v_n^1) \, \mathrm{d}x \, \mathrm{d}\tau \, .$$

We have performed an integration in time, and used $v_{0n}^1 \ge v_{0n}^2$. But, since $f^1 \ge f^2$,

$$\begin{split} [f^2(v_n^2) - f^1(v_n^1)] \operatorname{sign}^+(v_n^2 - v_n^1) &= [f^2(v_n^2) - f^2(v_n^1)] \operatorname{sign}^+(v_n^2 - v_n^1) \\ &+ [f^2(v_n^1) - f^1(v_n^1)] \operatorname{sign}^+(v_n^2 - v_n^1) \leq \mu (v_n^2 - v_n^1)_+ \,. \end{split}$$

Therefore

$$\int_{\Omega} (v_n^2 - v_n^1)_+(x, t) \, \mathrm{d}x \le \mu \iint_{Q_t} (v_n^2 - v_n^1)_+ \, \mathrm{d}x \, \mathrm{d}\tau \, .$$

Finally, an application of Gronwall's lemma concludes the proof.

CHAPTER 3

The relaxed formulation

3.1. A relaxed Stefan problem

Assume that in a material subject to a change of phase the enthalpy density is given by (in non-dimensional form)

$$v(x,t) = u(x,t) + \psi(x,t),$$
 (3.1)

where u is the temperature, and ψ is the concentration of the more energetic phase. The temperature of change of phase is u = 0. Therefore

$$\begin{split} \psi(x,t) &= 0\,, & u(x,t) < 0\,, \\ \psi(x,t) &\in [0,1]\,, & u(x,t) = 0\,, \\ \psi(x,t) &= 1\,, & u(x,t) > 0\,. \end{split}$$

In other words

$$\psi \in H(u)$$
, i.e., $\eta(\psi) \ni u$, (3.2)

where ${\cal H}$ is Heaviside's graph

$$H(s) = \begin{cases} s, & s < 0, \\ [0,1], & s = 0, \\ s+1, & s > 0, \end{cases}$$

and η its inverse

$$\eta(s) = \begin{cases} (-\infty, 0], & s = -1, \\ 0, & 0 < s < 1, \\ [0, \infty), & s = 1. \end{cases}$$

Note that (3.2) implies that the two phases can coexist only at the critical temperature u = 0; if $u(x, t) \neq 0$ only one pure phase may exist at (x, t). In this sense (3.2) is an equilibrium condition. Specifically, it does not allow for any superheating or undercooling effects (defined respectively as $\psi < 1$ at some points where u > 0, and $\psi > 0$ at some points where u < 0). Instead of (3.2) we can prescribe the non-equilibrium relation

$$\frac{\partial \psi}{\partial t} + \eta(\psi) \ni mu. \tag{3.3}$$

Here m > 0 is a parameter whose significance will be discussed below. The precise mathematical meaning of condition (3.3) is clarified in Appendix D; we recall that as a consequence of (3.3) we have

$$0 \le \psi \le 1 \,, \tag{3.4}$$

$$\frac{\partial \psi}{\partial t} = mu$$
, where $0 < \psi < 1.$ (3.5)

3.1.1. Formulation of the problem. Equation (3.3) must be completed with the initial data

$$\psi(x,0) = \psi_0, \qquad x \in \Omega, \tag{3.6}$$

where

$$\psi_0 \in L^{\infty}(\Omega), \qquad 0 \le \psi_0 \le 1.$$
(3.7)

Moreover the non-dimensional temperature must satisfy the parabolic problem

$$u_t + \psi_t - \Delta u = 0, \qquad \text{in } Q_T, \qquad (3.8)$$

$$\frac{\partial u}{\partial \nu} = 0, \qquad \text{on } \partial \Omega \times (0, T), \qquad (3.9)$$

$$u(x,0) = u_0(x), \qquad x \in \Omega.$$
 (3.10)

We could allow for a volumetric source term $f \in L^2(Q_T)$ on the right hand side of (3.8): we assume the heat equation to be homogeneous for the sake of brevity.

Definition 3.1. Let $u_0 \in L^{\infty}(\Omega)$, and (3.7) be satisfied. A couple $(u, \psi) \in L^{\infty}(Q_T)^2$ is a solution to problem (3.3), (3.6), (3.8)–(3.10) if

$$u \in L^2(0, T; W_1^2(\Omega)), \qquad \psi \in W_1^2(0, T; L^2(\Omega)),$$
 (3.11)

and for all $\varphi \in W_1^2(Q_T)$ we have

$$\iint_{Q_T} \{-(u+\psi)\varphi_t + \nabla u \cdot \nabla \varphi\} \,\mathrm{d}x \,\mathrm{d}t = \int_{\Omega} (u_0 + \psi_0)\varphi(x,0) \,\mathrm{d}x \,. \tag{3.12}$$

In addition we prescribe that

$$\iint_{Q_T} \left(\frac{\partial \psi}{\partial t}(x,t) - mu(x,t) \right) (\zeta(x,t) - \psi(x,t)) \, \mathrm{d}x \, \mathrm{d}t \ge 0 \tag{3.13}$$

for all $\zeta \in L^{\infty}(Q_T), 0 \leq \zeta \leq 1$.

For further information on the subject of this Chapter we refer the reader to [20] and to the bibliography therein.

3.2. Existence of solutions to the relaxed problem

We consider the problem with delay obtained from the one in Definition 3.1 by replacing u(x,t) with u(x,t-1/n), $n \in \mathbb{N}$ in (3.13). If t < 1/n we assume here $u(x,t-1/n) = u_0(x)$. This problem has a solution since, dividing the interval (0, T + 1/n) in subintervals of length 1/n, we may first solve the variational inequality assuming that u in it is known, and then solve the heat equation for the resulting function ψ . Let us denote by (u_n, ψ_n) such a solution.

Next we derive some integral estimates for (u_n, ψ_n) which are uniform on n. Routine calculations yield for 0 < t < T

$$\int_{\Omega(t)} u_n^2 \,\mathrm{d}x + \int_0^t \int_{\Omega} |\nabla u_n|^2 \,\mathrm{d}x \,\mathrm{d}\tau \le \int_0^t \int_{\Omega} (\psi_{nt}^2 + u_n^2) \,\mathrm{d}x \,\mathrm{d}\tau \,. \tag{3.14}$$

From (D.16) we obtain on the other hand

$$\iint_{Q_t} \psi_{n\tau}^2 \,\mathrm{d}x \,\mathrm{d}\tau \le \iint_{Q_t} m^2 u_n^2 \,\mathrm{d}x \,\mathrm{d}\tau + \frac{1}{n} \int_{\Omega} m^2 u_0^2 \,\mathrm{d}x \,. \tag{3.15}$$

Collecting (3.14) and (3.15), we obtain from Gronwall's lemma

$$\sup_{0 < \tau < t} \int_{\Omega(\tau)} u_n^2 \, \mathrm{d}x + \iint_{Q_t} |\nabla u_n|^2 \, \mathrm{d}x \, \mathrm{d}\tau$$

$$\leq \gamma \left(e^{2t} + \frac{m^2}{n} (e^{2t} - 1) \right) \int_{\Omega} u_0^2 \, \mathrm{d}x \,, \qquad 0 < t < T \,. \quad (3.16)$$

Hence, again by (3.15),

$$\iint_{Q_T} \psi_{n\tau}^2 \,\mathrm{d}x \,\mathrm{d}\tau \le \gamma \big(m, T, \|u_0\|_{2,\Omega}\big) \,, \tag{3.17}$$

where γ does not depend on n. Next, for any fixed $0 < \bar{t} < t < T$ we use as a testing function in (3.12)

$$u_{nt}\zeta(t)$$
, $\zeta(t) = \max\left(0, \min\left((t-\bar{t}/2)2\bar{t}^{-1}, 1\right)\right);$

note that this step in general requires a regularization procedure, since u_{nt} does not in general have the required smoothness. Formally we proceed as follows, by repeatedly integrating by parts

$$-\int_{0}^{t}\int_{\Omega} \{u_{n\tau}^{2}\zeta - u_{n\tau}\zeta\psi_{n\tau}\} \,\mathrm{d}x \,\mathrm{d}\tau$$
$$=\int_{0}^{t}\int_{\Omega} \{\nabla u_{n} \cdot \nabla u_{n\tau}\zeta\} \,\mathrm{d}x \,\mathrm{d}\tau = \frac{1}{2}\int_{\Omega(t)} |\nabla u_{n}|^{2}\zeta \,\mathrm{d}x - \frac{1}{2}\int_{0}^{t}\int_{\Omega} |\nabla u_{n}|^{2}\zeta' \,\mathrm{d}x \,\mathrm{d}\tau.$$

Using (3.16), (3.17), it follows from an application of Cauchy-Schwarz inequality that

$$\int_{\bar{t}}^{T} \int_{\Omega} u_{n\tau}^{2} \,\mathrm{d}x \,\mathrm{d}\tau + \int_{\Omega(t)} |\nabla u_{n}|^{2} \,\mathrm{d}x \leq \gamma \left(m, T, \|u_{0}\|_{2,\Omega}, \bar{t}\right), \qquad (3.18)$$

for all $0 < \overline{t} < t < T$.

As a consequence of (3.16), (3.17), (3.18) we may assume, perhaps by extracting a subsequence,

$$\psi_n, u_n \to \psi, u, \qquad \text{in } L^2(Q_T), \text{ and a.e. in } Q_T; \qquad (3.19)$$

$$\psi_{nt}, \nabla u_n \to \psi_t, \nabla u, \quad \text{weakly in } L^2(Q_T).$$
 (3.20)

Note that in (3.19) to prove the a.e. convergence of ψ_n we need a suitable equicontinuity in the $L^2(Q_T)$ norm which in turn follows from Proposition D.20, where we take

$$\psi_1(t) = \psi_n(x,t), \quad \psi_2(t) = \psi_n(x+h,t)$$

$$f_1(x,t) = u_n(x,t-1/n), \quad f_2(x,t) = u_n(x+h,t-1/n),$$

for a (small) $h \in \mathbf{R}^N$.

The fact that (u, ψ) is a solution in the sense of Definition 3.1 follows by taking the limit in (3.12), (3.13). While the first limit is trivial, the quantity in the approximating version of (3.13) amounts to

$$\iint_{Q_T} \left(\frac{\partial \psi_n}{\partial t}(x,t) - m u_n(x,t-1/n) \right) (\zeta(x,t) - \psi_n(x,t)) \, \mathrm{d}x \, \mathrm{d}t \ge 0 \,,$$

and we must only check that, setting $\tilde{u}_n(x,t) = u_n(x,t-1/n)$,

$$\tilde{u}_n \to u, \quad \text{in } L^2(Q_T).$$
(3.21)

However for any fixed $T > \overline{t} > 0$ we have from (3.16), (3.18) that

$$\iint_{Q_T} |u_n - \tilde{u}_n|^2 \, \mathrm{d}x \, \mathrm{d}t \le \frac{T}{n} \int_{\bar{t}}^T \int_{\Omega} u_{nt}^2 \, \mathrm{d}x \, \mathrm{d}t + \gamma \bar{t} [\|u_0\|_{2,\Omega}^2 + \sup_{0 < t < \bar{t}} \|u(\cdot, t)\|_{2,\Omega}^2] \\ \le \gamma (m, T, \|u_0\|_{2,\Omega}, \bar{t}) \frac{1}{n} + \gamma (m, T, \|u_0\|_{2,\Omega}) \bar{t} \, .$$

This shows that indeed $u_n - \tilde{u}_n \to 0$ in $L^2(Q_T)$.

Theorem 3.2. A solution in the sense of Definition 3.1 exists and is unique. More generally, if (u_1, ψ_1) , (u_2, ψ_2) are two solutions corresponding to initial data u_{01} , ψ_{01} and u_{02} , ψ_{02} , then

$$\|u_1 - u_2\|_{2,Q_T} + \sup_{0 < t < T} \|\psi_1 - \psi_2\|_{2,\Omega}$$

$$\leq \gamma(m,T) \Big(\|u_{01} - u_{02}\|_{2,\Omega} + \|\psi_{01} - \psi_{02}\|_{2,\Omega} \Big).$$
 (3.22)

PROOF. The existence has been obtained above, when we invoke also Exercise 3.3.

Assume that (u_1, ψ_1) , (u_2, ψ_2) are two solutions, and define $\tilde{u} = u_1 - u_2$, $\tilde{\psi} = \psi_1 - \psi_2$,

$$\tilde{w}(x,t) = \int_{0}^{t} \tilde{u}(x,\tau) \,\mathrm{d} au$$

Essentially the same argument employed below to get (3.33) leads us to

$$\int_{0}^{t} \int_{\Omega} \tilde{u}^{2} dx d\tau + \frac{1}{2} \int_{\Omega} |\nabla \tilde{w}(x,t)|^{2} dx =$$
$$\int_{\Omega} (\tilde{u}_{0} + \tilde{\psi}_{0}) \tilde{w} dx d\tau + \int_{0}^{t} \int_{\Omega} \tilde{\psi} \tilde{u} dx d\tau, \quad (3.23)$$

where $\tilde{u}_0 = u_{01} - u_{02}$, $\tilde{\psi}_0 = \psi_{01} - \psi_{02}$. Moreover we have, as in (D.21),

$$\int_{\Omega} \tilde{\psi}(x,t)^2 \, \mathrm{d}x \le 2m \int_{0}^{\bullet} \int_{\Omega} \tilde{u}\tilde{\psi} \, \mathrm{d}x \, \mathrm{d}\tau + \int_{\Omega} \tilde{\psi}_0(x)^2 \, \mathrm{d}x \,. \tag{3.24}$$

Adding (3.23) to (3.24) we infer, after applying Cauchy-Schwarz inequality,

$$\begin{split} \int_{0}^{\circ} \int_{\Omega} \tilde{u}^2 \, \mathrm{d}x \, \mathrm{d}\tau + \frac{1}{2} \int_{\Omega} |\nabla \, \tilde{w}(x,t)|^2 \, \mathrm{d}x + \int_{\Omega} \tilde{\psi}(x,t)^2 \, \mathrm{d}x \\ & \leq \gamma(T) \int_{\Omega} (\tilde{u}_0^2 + \tilde{\psi}_0^2) \, \mathrm{d}x + \gamma \int_{0}^{t} \int_{\Omega} \tilde{\psi}^2 \, \mathrm{d}x \, \mathrm{d}\tau \, . \end{split}$$

Finally the statement follows from Gronwall's lemma.

3.2.1. Exercises.

4

Exercise 3.3. Prove that $u_n \in L^{\infty}(Q_T)$ and that

$$||u||_{\infty,Q_T} \le ||u_0||_{\infty,Q_T}.$$

[See Section (A.4).]

3.3. The Stefan problem as limit of the relaxed problem

Here we show that as $m \to \infty$ the solution to the relaxed Stefan problem defined in Section 3.1 approaches the solution to the standard Stefan problem which was introduced in Chapter 2.

Assume that $m \in \mathbf{N}$ and denote by (u_m, ψ_m) the corresponding solution in the sense of Definition 3.1. Since we want to perform the limit as $m \to \infty$, we are interested in estimates for (u_m, ψ_m) which are uniform on m. It can be readily checked that the arguments in Section 3.2 yield

$$\sup_{0<\tau
$$\leq \gamma(T) \int_{\Omega} u_0^2 \,\mathrm{d}x \,. \quad (3.25)$$$$

In addition of course we have

$$0 \le \psi_m \le 1. \tag{3.26}$$

Unfortunately these bounds only allow us to infer weak convergence, up to extracting a subsequence, i.e.,

 $u_m, \nabla u_m \to u, \nabla u, \quad \text{weakly in } L^2(Q_T);$ (3.27)

$$\psi_m \to \psi$$
, weakly in $L^2(Q_T)$. (3.28)

This is enough to take the limit in (3.12); setting

$$v=u+\psi\,,$$

we can see, going back to Definition 2.1, that (2.14) is therefore satisfied, as well as the other assumptions on v, provided we are able to show that

$$\psi \in H(u)$$
, a.e. in Q_T . (3.29)

One additional piece of information we may directly extract from (3.25) is

$$\iint_{Q_T} \psi_{mt}^2 \,\mathrm{d}x \,\mathrm{d}t \le \gamma m \,. \tag{3.30}$$

Thus ψ_{mt}/m goes to zero strongly in $L^2(Q_T)$, so that, if also u_m converged strongly, we could take the limit in (3.13), or more exactly in the inequality obtained by dividing (3.13) by m, and get

$$\iint_{Q_T} u(x,t)(\psi(x,t) - \zeta(x,t)) \,\mathrm{d}x \,\mathrm{d}t \ge 0 \tag{3.31}$$

for all $\zeta \in L^{\infty}(Q_T)$, $0 \leq \zeta \leq 1$, whence (3.29) would follow.

3.3.1. Exploiting monotonicity. However, due to the weak character of our compactness results this approach is unfeasible. We resort to the following more complex argument based on monotonicity. First introduce the functions

$$w_m(x,t) = \int_0^t u_m(x,\tau) \,\mathrm{d}\tau, \qquad w(x,t) = \int_0^t u(x,\tau) \,\mathrm{d}\tau.$$

Note that for each 0 < t < T

$$\|w_m(\cdot,t)\|_{2,\Omega} \le \gamma \|u_m\|_{2,Q_T}, \qquad \|\nabla w_m(\cdot,t)\|_{2,\Omega} \le \gamma \|\nabla u_m\|_{2,Q_T}.$$

Moreover owing to (3.27) for each 0 < t < T

$$w_m(\cdot, t), \nabla w_m(\cdot, t) \to w(\cdot, t), \nabla w(\cdot, t), \quad \text{weakly in } L^2(\Omega).$$
 (3.32)

Proposition 3.4. The relation (3.29) holds true.

PROOF. Take as a testing function in (3.12)

$$\varphi(x,t) = \int_{t}^{T} u_m(x,\tau) \,\mathrm{d}\tau$$

We get

$$\iint_{Q_T} (u_m + \psi_m) u_m \, \mathrm{d}x \, \mathrm{d}\tau - \int_{\Omega} (u_0 + \psi_0) \varphi(x, 0) \, \mathrm{d}x = -\iint_{Q_T} \nabla \, u_m \cdot \nabla \, \varphi \, \mathrm{d}x \, \mathrm{d}\tau$$
$$= \iint_{Q_T} \frac{\partial}{\partial \tau} \frac{1}{2} \left| \nabla \, \varphi \right|^2 \, \mathrm{d}x \, \mathrm{d}\tau = -\frac{1}{2} \int_{\Omega} \left| \nabla \, w_m(x, T) \right|^2 \, \mathrm{d}x \,. \quad (3.33)$$

Therefore, by invoking the basic result of weak lower semicontinuity of the L^2 norm and (3.32) we get

$$\limsup_{m \to \infty} \iint_{Q_T} \psi_m u_m \, \mathrm{d}x \, \mathrm{d}\tau$$

$$\leq \int_{\Omega} (u_0 + \psi_0) w(x, T) \, \mathrm{d}x - \frac{1}{2} \int_{\Omega} |\nabla w(x, T)|^2 \, \mathrm{d}x - \iint_{Q_T} u^2 \, \mathrm{d}x \, \mathrm{d}\tau \,. \quad (3.34)$$

On the other hand, reasoning as in Subsection 2.4.1 we can prove starting from (2.14), that for a.e. 0 < t < T

+

$$\int_{\Omega(t)} (u+\psi)\zeta(x) \,\mathrm{d}x + \int_{0}^{t} \int_{\Omega} \nabla u \cdot \nabla \zeta \,\mathrm{d}x \,\mathrm{d}\tau = \int_{\Omega} (u_0+\psi_0)\zeta(x) \,\mathrm{d}x,$$

for all $\zeta \in W_1^2(\Omega)$. On selecting above $\zeta = u(\cdot, t)$ and integrating over (0, T)we find

$$\iint_{Q_T} (u+\psi)u\,\mathrm{d}x\,\mathrm{d}t = -\iint_{Q_T} \nabla w \cdot \nabla u\,\mathrm{d}x\,\mathrm{d}t + \int_{\Omega} (u_0+\psi_0)w(x,T)\,\mathrm{d}x$$
$$= -\frac{1}{2}\int_{\Omega} |\nabla w(x,T)|^2\,\mathrm{d}x + \int_{\Omega} (u_0+\psi_0)w(x,T)\,\mathrm{d}x. \quad (3.35)$$

From (3.34) and (3.35) we infer

$$\limsup_{m \to \infty} \iint_{Q_T} \psi_m u_m \, \mathrm{d}x \, \mathrm{d}\tau \le \iint_{Q_T} \psi_u \, \mathrm{d}x \, \mathrm{d}t \,. \tag{3.36}$$

Thus for any fixed ζ as in Definition 3.1 and for a suitable subsequence $\{m'\}$ we have

$$\begin{split} \iint_{Q_T} (\psi - \zeta) u \, \mathrm{d}x \, \mathrm{d}t &= \iint_{Q_T} \psi u \, \mathrm{d}x \, \mathrm{d}t - \iint_{Q_T} \zeta u \, \mathrm{d}x \, \mathrm{d}t \\ &\geq \limsup_{m \to \infty} \iint_{Q_T} \psi_m u_m \, \mathrm{d}x \, \mathrm{d}t - \lim_{m \to \infty} \iint_{Q_T} \zeta u_m \, \mathrm{d}x \, \mathrm{d}t \\ &= \lim_{m' \to \infty} \iint_{Q_T} (\psi_{m'} - \zeta) u_{m'} \, \mathrm{d}x \, \mathrm{d}t \\ &\geq \lim_{m' \to \infty} \frac{1}{m'} \iint_{Q_T} (\psi_{m'} - \zeta) \psi_{m't} \, \mathrm{d}x \, \mathrm{d}t = 0 \,, \end{split}$$

by virtue of the already remarked convergence $\psi_{mt}/m \to 0$ in $L^2(Q_T)$. This amounts to (3.31), whence the statement.

3.3.2. Exercises.

Exercise 3.5. Prove that (3.29) follows from (3.31).

•

APPENDIX A

Maximum principles for parabolic equations

A.1. The weak maximum principle

Let Q_T be a bounded open set of \mathbf{R}^{N+1} , contained in $\mathbf{R}^N \times (0,T)$, where T > 0.

Definition A.1. We denote by Q_T^* the parabolic interior of Q_T , that is the set of all points (\bar{x}, \bar{t}) with the property

$$\exists \varepsilon > 0: \quad B_{\varepsilon}(\bar{x}, \bar{t}) \cap \{t < \bar{t}\} \subset Q_T.$$

Here $B_{\varepsilon}(\bar{x}, \bar{t})$ denotes the (N+1)-dimensional ball with radius ε and center (\bar{x}, \bar{t}) . Define also the parabolic boundary $\partial_p Q_T$ of Q_T , as

$$\partial_p Q_T = \overline{Q_T} - Q_T^* \,.$$

The set $\partial_p Q_T$ is the parabolic analogue of the boundary of Q_T , i.e., roughly speaking, the region where initial and boundary data should be prescribed for parabolic problems set in Q_T (see Figure A.1). For example, if $Q_T = (0, L) \times (0, T)$ then $Q_T^* = (0, L) \times (0, T]$. Obviously we have $Q_T \subset Q_T^* \subset \overline{Q_T}$. In general $\partial_p Q_T$ is not a closed set.



FIGURE A.1. The dashed lines and the point E belong to the parabolic interior, but the points A, B, C, D, as well as the solid lines, belong to the parabolic boundary.

In the following we denote

$$\begin{aligned} \mathcal{L}u &= u_t - a_{ij}(x,t)u_{x_ix_j} + b_i(x,t)u_{x_i} + c(x,t)u \,, \\ \mathcal{L}_0 u &= u_t - a_{ij}(x,t)u_{x_ix_j} + b_i(x,t)u_{x_i} \,. \end{aligned}$$

Throughout this Appendix we employ the summation convention, and assume that

$$u \in C^{2,1}(Q_T^*) \cap C(\overline{Q_T}), \qquad a_{ij}, b_i, c \in C(Q_T^*);$$
$$a_{ij}(x,t)\xi_i\xi_j \ge \nu|\xi|^2, \qquad \text{for all } \xi \in \mathbf{R}^N, (x,t) \in Q_T^*$$

Here $\nu > 0$ is a given constant. We also assume

$$\sum_{i,j=1}^{N} \|a_{ij}\|_{\infty} =: A < \infty, \quad \sum_{i=1}^{N} \|b_i\|_{\infty} =: B < \infty, \quad \|c\|_{\infty} =: C < \infty.$$

Lemma A.2. Assume that $\mathcal{L}_0 u(\bar{x}, \bar{t}) < 0$, where $(\bar{x}, \bar{t}) \in Q_T^*$. Then u can not attain a local maximum at (\bar{x}, \bar{t}) .

PROOF. Recalling the definition of Q_T^* , we have, reasoning by contradiction,

$$\mathcal{L}_0 u(\bar{x}, \bar{t}) = u_t(\bar{x}, \bar{t}) - a_{ij}(\bar{x}, \bar{t}) u_{x_i x_j}(\bar{x}, \bar{t}) + b_i(\bar{x}, \bar{t}) u_{x_i}(\bar{x}, \bar{t})$$
$$= u_t(\bar{x}, \bar{t}) - a_{ij}(\bar{x}, \bar{t}) u_{x_i x_j}(\bar{x}, \bar{t}) .$$

Note that, if (\bar{x}, \bar{t}) is a point of local maximum, then $u_t(\bar{x}, \bar{t}) \ge 0$. By the same token, $a_{ij}u_{x_ix_j} \le 0$ at (\bar{x}, \bar{t}) , as we show below. This leads of course to an inconsistency, as $\mathcal{L}_0u(\bar{x}, \bar{t}) < 0$ by assumption.

To prove the assertion $a_{ij}u_{x_ix_j} \leq 0$ at (\bar{x}, \bar{t}) , we change spatial coordinates defining $y = \bar{x} + \Gamma(x - \bar{x})$, and v(y, t) = u(x(y), t), where $\Gamma = (\gamma_{ij})$ is an $N \times N$ matrix such that $\Gamma(a_{ij}(\bar{x}, \bar{t}))\Gamma^t$ coincides with the diagonal matrix diag $(\lambda_1, \ldots, \lambda_N)$ (we may assume without loss of generality that (a_{ij}) is symmetric). Then

$$a_{ij}u_{x_ix_j} = a_{ij}v_{y_hy_k}\gamma_{hi}\gamma_{kj} = \lambda_h v_{y_hy_h}, \quad \text{at } (\bar{x}, t).$$

Note that $v(\cdot, \bar{t})$ attains a local maximum at $y = \bar{x}$, so that $v_{y_h y_h} \leq 0$ for all h. We also take into account that $\lambda_h > 0$ for all h, since (a_{ij}) is positive definite. The result immediately follows.

Theorem A.3. (WEAK MAXIMUM PRINCIPLE) Let $\mathcal{L}_0 u \leq 0$ in Q_T^* . Then

$$\max_{\overline{Q_T}} u = \sup_{\partial_p Q_T} u. \tag{A.1}$$

PROOF. Let us define

Į

$$v = (u - M)e^{-\varepsilon t}$$
, $M = \sup_{\partial_p Q_T} u$, $\varepsilon > 0$.

If u - M is positive at some $(x, t) \in Q_T^*$, then v attains a positive maximum somewhere in Q_T^* , say at (\bar{x}, \bar{t}) . Indeed $v \leq 0$ on $\partial_p Q_T$. We calculate

$$v_t = u_t e^{-\varepsilon t} - \varepsilon v$$
, $v_{x_i} = u_{x_i} e^{-\varepsilon t}$, $v_{x_i x_j} = u_{x_i x_j} e^{-\varepsilon t}$.

Therefore

$$\mathcal{L}_0 v(\bar{x}, \bar{t}) = e^{-\varepsilon t} \mathcal{L}_0 u(\bar{x}, \bar{t}) - \varepsilon v(\bar{x}, \bar{t}) \le -\varepsilon v(\bar{x}, \bar{t}) < 0.$$

Upon recalling Lemma A.2, this inconsistency concludes the proof.

A.1.1. More general operators. Let us consider here more general operators of the form \mathcal{L} .

Theorem A.4. Let $\mathcal{L}u \leq f$ in Q_T^* , $f \in C(\overline{Q_T})$. Define

$$\int_{0}^{t} \|f_{+}(\cdot,\tau)\|_{\infty} \,\mathrm{d}\tau =: H(t), \qquad 0 < t < T.$$
(A.2)

Then

$$\max_{\overline{Q_T}} u_+ \le e^{C_-T} \left(\sup_{\partial_p Q_T} u_+ + H(T) \right), \tag{A.3}$$

where $C_{-} = \|c_{-}\|_{\infty,Q_{T}}$.

PROOF. Define for a constant $\gamma > 0$ to be chosen

$$w(x,t) = e^{-\gamma t}u(x,t) - m - H(t),$$

where

$$m := \sup_{\partial_p Q_T} u_+ \, .$$

By definition of m, and since $\gamma > 0$, we have $w \leq 0$ on $\partial_p Q_T$. Moreover

$$\mathcal{L}_{0}w = e^{-\gamma t}\mathcal{L}_{0}u - \gamma e^{-\gamma t}u - \|f_{+}(\cdot,t)\|_{\infty}$$

$$\leq -ce^{-\gamma t}u + e^{-\gamma t}f_{+}(x,t) - \gamma e^{-\gamma t}u - \|f_{+}(\cdot,t)\|_{\infty} \leq -[c+\gamma]e^{-\gamma t}u.$$

Therefore we have $\mathcal{L}_0 w < 0$ where w > 0 (and hence u > 0 too), provided we select $\gamma = C_- + \varepsilon$ for any arbitrarily fixed $\varepsilon > 0$. Thus, if w attains a positive maximum in Q_T^* , we arrive at an inconsistency with Lemma A.2. We conclude that $w \leq 0$ in $\overline{Q_T}$, and we recover (A.3) on letting $\varepsilon \to 0$. \Box

Note that according to our theorem above, if $u \leq 0$ on $\partial_p Q_T$, and $f \leq 0$ in Q_T , then $u \leq 0$ in Q_T , regardless of the sign of c.

Remark A.5. All the results of this section still hold if the parabolicity constant ν is equal to 0. This is not the case for the results in next two sections. See also Section A.4.

A.2. The strong maximum principle

Let $(\bar{x}, \bar{t}) \in Q_T^*$. Define the set $\mathcal{S}(\bar{x}, \bar{t})$ as the set of all $(x, t) \in Q_T$ with the property

(x,t) can be connected to (\bar{x},\bar{t}) by a polygonal contained in Q_T ,

along which t is increasing, when going from (x, t) to (\bar{x}, \bar{t}) .

A polygonal is a connected curve made of a finite number of straight line segments. Essentially, the strong maximum principle asserts that if $\mathcal{L}_0 u \leq 0$ in Q_T , and u attains its maximum at a point (\bar{x}, \bar{t}) of Q_T^* , then u is constant in $\mathcal{S}(\bar{x}, \bar{t})$. Our first result is a weaker version of this principle.

The proof we present here was taken from $[2]^1$. The strong maximum principle for parabolic equations was first proven in [13].

¹The author of [2] quotes a course of D. Aronson (Minneapolis) as source of the proof.



FIGURE A.2. The set $S(\bar{x}, \bar{t})$ as defined in the text.

Lemma A.6. Let $\mathcal{L}_0 v \leq 0$ in $P_T = B_{\delta}(0) \times (0,T)$, $v \in C^{2,1}(P_T^*) \cap C(\overline{P_T})$, where $\delta > 0$, T > 0 and we assume

$$v(x,t) \le M$$
, $|x| = \delta$, $0 < t < T$, (A.4)

$$v(x,0) < M \,, \qquad |x| \le \delta \,, \tag{A.5}$$

for a given constant M. Then

$$v(x,T) < M, \qquad |x| < \delta. \tag{A.6}$$

PROOF. We have by continuity

$$v(x,0) < M - \varepsilon \delta^4$$
, $|x| \le \delta$, (A.7)

for a suitable $\varepsilon > 0$, which we fix from now on subject to this constraint. Let us define, for a $\alpha > 0$ to be chosen,

$$w(x,t) = M - \varepsilon(\delta^2 - |x|^2)^2 e^{-\alpha t} - v(x,t).$$

Then

$$\mathcal{L}_{0}w(x,t) = -\mathcal{L}_{0}[\varepsilon(\delta^{2} - |x|^{2})^{2}e^{-\alpha t}] - \mathcal{L}_{0}v(x,t)$$

$$\geq -\mathcal{L}_{0}[\varepsilon(\delta^{2} - |x|^{2})^{2}e^{-\alpha t}] = e^{-\alpha t} \left\{ \varepsilon \alpha (\delta^{2} - |x|^{2})^{2} - 4\varepsilon (\delta^{2} - |x|^{2})a_{jj} + 8\varepsilon a_{ij}x_{i}x_{j} - 4\varepsilon b_{i}x_{i}(\delta^{2} - |x|^{2}) \right\}.$$
 (A.8)

Here we employ the summation convention even for the term a_{jj} and we understand the coefficient a_{ij} , b_i to be calculated at (x, t). We aim at proving that the quantity $\{\ldots\}$ in last formula above is non negative, for a suitable choice of $\alpha > 0$. Introduce a parameter $\tau \in (0, 1)$, and distinguish the cases:

(i) If $|x| \leq \tau \delta$, then

$$\{\dots\} \ge \varepsilon (\delta^2 - |x|^2) \left[\alpha \delta^2 (1 - \tau^2) - 4A - 4B\tau \delta \right] \ge 0,$$

provided

$$\alpha \delta^2 (1 - \tau^2) - 4A - 4B\tau \delta \ge 0.$$
(A.9)

(ii) If
$$\delta > |x| > \tau \delta$$
, then
 $\{\dots\} \ge \varepsilon \left[-4A\delta^2(1-\tau^2) + 8\nu |x|^2 - 4B\delta^3(1-\tau^2) \right]$
 $\ge 4\varepsilon \left[-A\delta^2(1-\tau^2) + 2\nu\tau^2\delta^2 - B\delta^3(1-\tau^2) \right]$
 $= 4\varepsilon\delta^2 \left[2\nu\tau^2 - A(1-\tau^2) - B\delta(1-\tau^2) \right] \ge 0$,

provided

$$2\nu\tau^2 - A(1-\tau^2) - B\delta(1-\tau^2) \ge 0.$$
 (A.10)

We may first select τ so as (A.10) is satisfied, and then choose α so that (A.9) is satisfied too. Having fixed in this fashion the values of τ and α , we proceed to observe that

$$\mathcal{L}_0 w \ge 0, \qquad \text{in } P_T$$

Moreover on the parabolic boundary of P_T we have

$$w(x,t) = M - v(x,t) \ge 0, \qquad \text{on } |x| = \delta;$$

$$w(x,0) = M - \varepsilon(\delta^2 - |x|^2)^2 - v(x,0)$$

$$\ge M - \varepsilon\delta^4 - v(x,0) \ge 0, \qquad \text{in } |x| \le \delta;$$

we have made use of (A.7). Therefore $w \ge 0$ in P_T owing to the weak maximum principle.

Especially

$$w(x,T) = M - \varepsilon(\delta^2 - |x|^2)^2 e^{-\alpha T} - v(x,T) \ge 0,$$

and we finally prove our claim, i.e., for $|x| < \delta$,

$$v(x,T) \le M - \varepsilon (\delta^2 - |x|^2)^2 e^{-\alpha T} < M.$$

Theorem A.7. (STRONG MAXIMUM PRINCIPLE) Let $\mathcal{L}_0 u \leq 0$ in Q_T^* . If $(\bar{x}, \bar{t}) \in Q_T^*$, and

$$\max_{\overline{Q_T}} u = u(\bar{x}, \bar{t}) \,,$$

then

$$u(x,t) = u(\bar{x},\bar{t}), \quad \text{for all } (x,t) \in \mathcal{S}(\bar{x},\bar{t}).$$
 (A.11)

PROOF. Let us proceed by contradiction. Assume a point $(x_1, t_1) \in \mathcal{S}(\bar{x}, \bar{t})$ exists such that $u(x_1, t_1) < u(\bar{x}, \bar{t}) =: M$, and consider a polygonal (which must exist by definition of $\mathcal{S}(\bar{x}, \bar{t})$)

$$\bigcup_{i=1}^{n} \{ (1-\lambda)(x_i, t_i) + \lambda(x_{i+1}, t_{i+1}) \mid \lambda \in [0, 1] \},\$$

where $(x_{n+1}, t_{n+1}) = (\bar{x}, \bar{t})$, and $t_i < t_{i+1}$, for $i = 1, \ldots n$. We are going to prove that

 $u(x_i, t_i) < M \Longrightarrow u(x_{i+1}, t_{i+1}) < M$,

which obviously leads us to the contradiction

$$u(\bar{x}, \bar{t}) = u(x_{n+1}, t_{n+1}) < M = u(\bar{x}, \bar{t}).$$

We may assume without loss of generality i = 1, i + 1 = 2. Let us switch to different space coordinates:

$$\xi^j = x^j - x_1^j - (x_2^j - x_1^j) \frac{t - t_1}{t_2 - t_1}, \qquad j = 1, \dots, N.$$

DANIELE ANDREUCCI

Thus

$$(x_1, t_1) \mapsto (0, t_1), \qquad (x_2, t_2) \mapsto (0, t_2).$$

Also define the function

$$v(\xi, t) = u(x(\xi, t), t) \,.$$

In the change to the new variables, the set Q_T is transformed to an open set which certainly contains the closure of the cylinder

$$E_{\delta} = \{ (\xi, t) \mid |\xi| < \delta \, , \, t_1 < t < t_2 \, \} \, ,$$

provided $\delta > 0$ is suitably chosen. Possibly redefining δ we may assume (by continuity)

$$v(\xi, t_1) < M, \qquad |\xi| \le \delta,$$

From now on, let δ be fixed in this way. We have in E_{δ}

$$\tilde{\mathcal{L}}_0 v(\xi, t) := v_t(\xi, t) - a_{ij}(x(\xi, t), t) v_{\xi_i \xi_j}(\xi, t) + \tilde{b}_i(\xi, t) v_{\xi_i}(\xi, t) \le 0.$$
(A.12)

where

$$\tilde{b}_i(\xi, t) = b_i(x(\xi, t), t) - \frac{x_2^i - x_1^i}{t_2 - t_1}$$

Note that \mathcal{L}_0 is an operator satisfying the same assumptions as \mathcal{L}_0 . More specifically

$$\sum_{i=1}^{n} \|\tilde{b}_i\|_{\infty} \le \tilde{B} := B + N \frac{|x_2 - x_1|}{(t_2 - t_1)}.$$

Finally,

$$v(\xi, t) \le M$$
, $|\xi| = \delta$, $t_1 < t < t_2$,

follows from the definition of M. Therefore we may apply Lemma A.6 to conclude that

$$u(x_2, t_2) = v(0, t_2) < M$$
,

as claimed.

A.2.1. More general operators. Lemma A.6 and Theorem A.7 still hold, if \mathcal{L}_0 in their statements is replaced with the more general operator \mathcal{L} , provided $c(x,t)M \geq 0$ in Q_T (here $M = u(\bar{x}, \bar{t})$ in Theorem A.7).

We sketch here the changes needed in the proof of the Lemma, the proof of the Theorem being essentially the same:

The calculation in (A.8) should be replaced with

$$\begin{split} \tilde{\mathcal{L}}w(\xi,t) &= \tilde{\mathcal{L}}[M - \varepsilon(\delta^2 - |\xi|^2)^2 e^{-\alpha(t-t_1)}] - \tilde{\mathcal{L}}v(\xi,t) \\ &\ge cM - \tilde{\mathcal{L}}[\varepsilon(\delta^2 - |\xi|^2)^2 e^{-\alpha(t-t_1)}] \ge -C\varepsilon(\delta^2 - |\xi|^2)^2 e^{-\alpha(t-t_1)} + \mathcal{F} \\ &= e^{-\alpha(t-t_1)} \left\{ \varepsilon(\alpha - C)(\delta^2 - |\xi|^2)^2 \\ &- 4\varepsilon(\delta^2 - |\xi|^2)a_{jj} + 8\varepsilon a_{ij}\xi_i\xi_j - \left[b_i - \frac{x_2^i - x_1^i}{t_2 - t_1}\right] 4\varepsilon\xi_i(\delta^2 - |\xi|^2) \right\}, \end{split}$$

where \mathcal{F} denotes the last term in the chain of inequalities in (A.8). Here we used the fact that $cM \geq 0$. It is clear that, additionally assuming e.g., $\alpha > 2C$, the proof can be continued as above, taking into account Theorem A.4.

A.3. Hopf's lemma (the boundary point principle)

Definition A.8. We say that a point $(\bar{x}, \bar{t}) \in \partial_p Q_T$ has the property of the spherical cap if there exists an open ball $B_r(x_0, t_0)$ such that

$$(\bar{x},\bar{t}) \in \partial B_r(x_0,t_0), \qquad B_r(x_0,t_0) \cap \{t < \bar{t}\} \subset Q_T,$$

with $x_0 \neq \bar{x}$.

In the following we denote by $C_r(\bar{x}, \bar{t})$ a cap $B_r(x_0, t_0) \cap \{t < \bar{t}\}$ as the one appearing in Definition A.8. Note that if (\bar{x}, \bar{t}) has the property of the spherical cap, then there exist infinitely many such caps.

Remark A.9. If (\bar{x}, \bar{t}) has the property of the spherical cap, the *N*-dimensional open set $G := Q_T^* \cap \{t = \bar{t}\}$ has the usual property of the sphere at \bar{x} . Indeed, it contains the *N*-dimensional sphere $B_r(x_0, t_0) \cap \{t = \bar{t}\}$, which however touches the boundary of G at \bar{x} .

On the other hand, examples can be easily given where $(\bar{x}, \bar{t}) \in \partial_p Q_T$ has the property of the spherical cap, but fails to have the property of the sphere; see Figures A.3 and A.4.



FIGURE A.3. Every point of $\partial_p Q_T^1 \cap \{t > 0\}$ has the spherical cap property. This fails for the points on the vertical edges of Q_{Θ}^2 .

A version of Hopf's lemma for parabolic equations was first proven in [19], [7].

Theorem A.10. (HOPF'S LEMMA) Let $\mathcal{L}_0 u \leq 0$ in Q_T^* . Let $(\bar{x}, \bar{t}) \in \partial_p Q_T$ have the property of the spherical cap. If

$$u(x,t) < u(\bar{x},\bar{t}), \quad \text{for all } (x,t) \in C_r(\bar{x},\bar{t}), \quad (A.13)$$

then

$$\frac{\partial u}{\partial \mathbf{e}}(\bar{x},\bar{t}) < 0, \qquad (A.14)$$

where $\mathbf{e} \in \mathbf{R}^{N+1}$ is any direction such that

$$(\bar{x}, \bar{t}) + s\mathbf{e} \in \overline{C_r(\bar{x}, \bar{t})}, \quad for \ 0 < s < \Sigma(\mathbf{e}),$$
 (A.15)

and we also assume that the derivative in (A.14) exists.

PROOF. First, let us invoke the strong maximum principle to prove that the maximum value $u(\bar{x}, \bar{t})$ may not be attained in the parabolic interior of $C_r(\bar{x}, \bar{t})$. Namely, we obtain in this fashion the additional piece of information that

$$u(x,t) < u(\bar{x},\bar{t}),$$
 for all $(x,\bar{t}) \in B_r(x_0,t_0).$

Then, for each fixed **e** as in (A.15), we may find a spherical cap C' such that its closure $\overline{C'}$ is contained in $C_r(\bar{x}, \bar{t})^* \cup \{(\bar{x}, \bar{t})\}$ and

$$u(x,\bar{t}) < u(\bar{x},\bar{t}), \quad \text{for all } (x,t) \in \overline{C'}, (x,t) \neq (\bar{x},\bar{t}), \quad (A.16)$$

$$(\bar{x}, \bar{t}) + s\mathbf{e} \in \overline{C'}, \quad \text{for } 0 < s < \Sigma'(\mathbf{e}).$$
 (A.17)

We'll keep the notation $C_r(\bar{x}, \bar{t})$ in the following for a cap satisfying (A.16), (A.17).

Let us consider the barrier function

$$w(x,t) = \exp\left\{-\alpha(|x-x_0|^2 + |t-t_0|^2)\right\} - \exp\{-\alpha r^2\},\$$

where $\alpha > 0$ is to be chosen, and (x_0, t_0) , r are as in Definition A.8. Thus 1 > w > 0 in $B_r(x_0, t_0)$, w = 0 on $\partial B_r(x_0, t_0)$, and

$$w_t(x,t) = -2(t-t_0)\alpha \exp\left\{-\alpha(|x-x_0|^2 + |t-t_0|^2)\right\},\$$

$$w_{x_i}(x,t) = -2(x^i - x_0^i)\alpha \exp\left\{-\alpha(|x-x_0|^2 + |t-t_0|^2)\right\},\$$

$$w_{x_ix_j}(x,t) = \left(-2\delta_{ij}\alpha + 4(x^i - x_0^i)(x^j - x_0^j)\alpha^2\right)\exp\{\dots\}.$$

Therefore we have

$$\mathcal{L}_{0}w(x,t) = 2\alpha \exp\{\dots\} \left[-(t-t_{0}) + a_{ii} - 2\alpha a_{ij}(x^{i} - x_{0}^{i})(x^{j} - x_{0}^{j}) - b_{i}(x^{i} - x_{0}^{i}) \right]$$

$$\leq 2\alpha \exp\{\dots\} \left[(t_{0} - t) + A - 2\alpha\nu|x - x_{0}|^{2} + B|x - x_{0}| \right]. \quad (A.18)$$

Define

$$\Omega = C_r(\bar{x}, \bar{t}) \cap \{ |x - \bar{x}| < \varepsilon \},\$$

where the positive number ε is selected so as, for $(x, t) \in \Omega$,

$$|x - x_0| \ge |\bar{x} - x_0| - |x - \bar{x}| \ge r \sin \theta - \varepsilon \ge \frac{1}{2} r \sin \theta.$$

Here θ is the angle between the *t* axis and the straight line joining (x_0, t_0) , (\bar{x}, \bar{t}) . Note that $\theta \in (0, \pi/2]$ as a consequence of $x_0 \neq \bar{x}$, according to Definition A.8; see also Figure A.4. Hence, in Ω we have

$$\mathcal{L}_0 w(x,t) \le 2\alpha \exp\{\dots\} \left[r + A - \frac{1}{2}\alpha\nu r^2 \sin^2\theta + Br\right] \le 0,$$

provided we finally choose α so that

$$\frac{2(A+r+Br)}{\nu r^2 \sin^2 \theta} \le \alpha \,.$$



FIGURE A.4. Ω and $C_r(\bar{x}, \bar{t})$, in the case $t_0 > \bar{t}$.

Define for a positive number μ to be chosen presently,

$$v(x,t) = u(x,t) + \mu w(x,t), \qquad (x,t) \in \Omega$$

Note that

 $\partial_p \Omega = S_1 \cup S_2$, $S_1 \subset \partial B_r(x_0, t_0)$, $S_2 \subset \{ |x - \bar{x}| = \varepsilon \} \cap \overline{C_r(\bar{x}, \bar{t})}$. Then, on $S_1 w = 0$ and thus

$$v(x,t) = u(x,t) \le u(\bar{x},\bar{t}).$$

On S_2 , taking into account (A.16),

$$u(x,t) \le u(\bar{x},\bar{t}) - \sigma$$

for a suitable $\sigma > 0$. It follows that on S_2 too

$$v(x,t) = u(x,t) + \mu w(x,t) \le u(\bar{x},\bar{t}) - \sigma + \mu \le u(\bar{x},\bar{t}),$$

if we choose $\mu \leq \sigma$. Moreover

$$\mathcal{L}_0 v = \mathcal{L}_0 u + \mu \mathcal{L}_0 w \le \mathcal{L}_0 u \le 0, \qquad \text{in } \Omega.$$

The weak maximum principle yields

$$v(x,t) \le u(\bar{x},\bar{t}), \quad \text{in } \Omega.$$

On the other hand, $v(\bar{x}, \bar{t}) = u(\bar{x}, \bar{t})$, so that

$$\frac{\partial v}{\partial \mathbf{e}}(\bar{x},\bar{t}) \le 0.$$

Therefore

$$\frac{\partial u}{\partial \mathbf{e}}(\bar{x},\bar{t}) = \frac{\partial v}{\partial \mathbf{e}}(\bar{x},\bar{t}) - \mu \frac{\partial w}{\partial \mathbf{e}}(\bar{x},\bar{t}) \le 2\mu\alpha(\bar{x}-x_0,\bar{t}-t_0) \cdot \mathbf{e}\exp\{\dots\} < 0.$$

DANIELE ANDREUCCI

A typical application of Theorem A.10 is the following: assume u satisfies $\mathcal{L}_0 u = 0$ in Q_T and attains its maximum at a point $(\bar{x}, \bar{t}) \in \partial_p Q_T$, having the spherical cap property. Unless u is identically constant in a portion of Q_T , by the strong maximum principle, u is strictly less than its maximum value in Q_T . Therefore we are in a position to apply Hopf's lemma, and prove $\frac{\partial u}{\partial \mathbf{n}} > 0$, if \mathbf{n} is the spatial outer normal to Q_T at (\bar{x}, \bar{t}) (as defined in Subsection 2.3.1 of Chapter 2).

A.3.1. More general operators. Theorem A.10 still holds, if \mathcal{L}_0 in the statement is replaced with the more general operator \mathcal{L} , provided $c(x,t) \geq 0$ in Q_T , and $u(\bar{x}, \bar{t}) \geq 0$. We sketch here the changes needed in the proof: The operator \mathcal{L}_0 is to be replaced everywhere with \mathcal{L} . Estimate (A.18) is substituted with the relation

$$\mathcal{L}w(x,t) \le 2\alpha \exp\{\dots\} \left[\frac{c}{2\alpha} - (t-t_0) + a_{ii} - 2\alpha a_{ij} (x^i - x_0^i) (x^j - x_0^j) - b_i (x^i - x_0^i) \right]$$

$$\le 2\alpha \exp\{\dots\} \left[A + \frac{C}{2} - \frac{1}{2}\alpha\nu r^2 \sin^2\theta + r + Br \right] \le 0,$$

which is valid in Ω as above, for a suitable selection of $\alpha > 1$. The proof is concluded as above.

A.3.2. Maximum estimates in problems with boundary conditions involving the spatial gradient. Assume *u* solves the problem, to be complemented with initial and additional boundary data,

$$\mathcal{L}_0 u = 0, \qquad \text{in } G \times (0, T), \tag{A.19}$$

$$\frac{\partial u}{\partial \mathbf{n}} = 0, \quad \text{on } \Gamma \subset \partial G \times (0, T),$$
 (A.20)

where $G \subset \mathbf{R}^N$, and **n** denotes the outer spatial normal to $\partial G \times (0, T)$. Then u can not attain either its maximum or its minimum on Γ , owing to Hopf's lemma (unless it is identically constant in some open set). Assume now (A.20) is replaced with

$$\frac{\partial u}{\partial \mathbf{n}} = -h(x,t)u + k(x,t), \quad \text{on } \Gamma \subset \partial G \times (0,T).$$

Here h > 0 and k are continuous functions. Assume $(\bar{x}, \bar{t}) \in \Gamma$ is a point of maximum for u. Then

$$0 < \frac{\partial u}{\partial \mathbf{n}}(\bar{x}, \bar{t}) = -h(\bar{x}, \bar{t})u(\bar{x}, \bar{t}) + k(\bar{x}, \bar{t}),$$

0

implying

$$u(\bar{x},\bar{t}) < \frac{k(\bar{x},\bar{t})}{h(\bar{x},\bar{t})}.$$

Note however that a similar, but not necessarily strict, inequality can be proven trivially without invoking Hopf's lemma. Analogous estimates hold at points of minimum.

A.4. Maximum principle for weak solutions

In this section we proceed formally, with the purpose of exhibiting the ideas behind a possible extension of the maximum principle to weak solutions of

$$u_t - \operatorname{div} \mathbf{a}(x, t, u, \nabla u) \le 0, \qquad \text{in } Q_T = G \times (0, T), \tag{A.21}$$

where

$$\mathbf{a}(x,t,u,\nabla u)\cdot\nabla u\geq 0\,.$$

Assume that $\partial G \times (0,T) = S_1 \cup S_2$, with

$$u(x,t) \le M, \quad \text{on } S_1,$$
$$\mathbf{a}(x,t,u,\nabla u) \cdot \mathbf{n} \le 0, \quad \text{on } S_2,$$

in a sense suitable for the calculations showed below, and also that

$$u(x,0) \le M \,, \qquad x \in G$$

Here \mathbf{n} is, as above, the outer spatial normal.

Multiply (formally) (A.21) by $(u - M)_+$, and integrate by parts, obtaining, on dropping the non negative term involving $\mathbf{a} \cdot \nabla u$,

$$\int_{G} (u(x,t) - M)_{+}^{2} dx \leq 2 \int_{0}^{t} \int_{\partial G} (u - M)_{+} \mathbf{a}(x,t,u,\nabla u) \cdot \mathbf{n} d\sigma d\tau + \int_{G} (u(x,0) - M)_{+}^{2} dx,$$

for all $t \in (0, T)$. The last integral equals 0, because of the assumed bound on the initial data. The surface integral is non positive: indeed, on S_1 we have $(u - M)_+ = 0$, while on S_2 it holds

$$(u-M)_+\mathbf{a}(x,t,u,\nabla u)\cdot\mathbf{n}\leq 0.$$

Therefore we get

$$\int_{G} (u(x,t) - M)_{+}^{2} \, \mathrm{d}x \le 0, \qquad \text{for all } 0 < t < T,$$

i.e., $u \leq M$ in $G \times (0, T)$.

APPENDIX B

A theorem by Kruzhkov

We present here a result of [10], which is instrumental in our proof of existence of weak solutions to the Stefan problem, in the modified version quoted in [8].

B.1. Mollifying kernels

Let φ be a mollifying kernel, i.e.,

$$\varphi\in C_0^\infty({\pmb R})\,,\quad {\rm supp}\,\varphi\subset [-1,1]\,,\quad \varphi\geq 0\,,\quad \varphi>0\quad {\rm on}\,\, [-1/2,1/2].$$
 Define

$$\varphi_{\varepsilon}(x) = \frac{1}{\varepsilon^N} \varphi(\frac{|x|}{\varepsilon}), \qquad x \in \mathbf{R}^N.$$

On multiplying, if required, φ by a positive constant, we may assume

$$\int_{\mathbf{R}^N} \varphi_{\varepsilon}(x) \, \mathrm{d}x = \int_{\mathbf{R}^N} \varphi(|x|) \, \mathrm{d}x = 1, \quad \text{for all } \varepsilon > 0.$$

Let $v \in L^1_{\text{loc}}(\mathbf{R}^N)$. Define for all $x \in \mathbf{R}^N$,

$$v_{\varepsilon}(x) = \int_{\mathbf{R}^N} v(y)\varphi_{\varepsilon}(x-y) \,\mathrm{d}y = \int_{\mathbf{R}^N} v(x-z)\varphi_{\varepsilon}(z) \,\mathrm{d}z \,. \tag{B.1}$$

Let Ω be a bounded open set of \mathbf{R}^N with a Lipschitz continuous boundary. Define for $1 > \delta > 0$

$$\Omega_{\delta} = \{ x \in \Omega \mid \operatorname{dist}(x, \partial \Omega) > \delta \}.$$

In the following ω_{δ} will denote a modulus of continuity, that is a continuous, non decreasing, non negative function $\omega_{\delta} : \mathbf{R} \to \mathbf{R}$ such that $\omega_{\delta}(0) = 0$. The notation emphasizes the possible dependence of this function on δ .

Our first result is not actually needed in the proof of the main estimate Theorem B.3, but it was quoted in the proof of Theorem 2.13 of Chapter 2.

Lemma B.1. Let $v \in L^1(\Omega)$, and assume that for each $0 < \delta < 1$, $h \in \mathbf{R}^N$, $|h| < \delta$

$$\int_{\Omega_{\delta}} |v(x+h) - v(x)| \, \mathrm{d}x \le \omega_{\delta}(|h|) \,. \tag{B.2}$$

Then, for all $0 < \varepsilon < \delta$, $|h| < \delta$,

$$\int_{\Omega_{2\delta}} |v_{\varepsilon}(x+h) - v_{\varepsilon}(x)| \, \mathrm{d}x \le \omega_{\delta}(|h|) \,. \tag{B.3}$$

Note that, to keep our notation formally coherent with (B.1), we extend in (B.3) v to v = 0 outside of Ω . However, the integrals defining $v_{\varepsilon}(x+h)$ and $v_{\varepsilon}(x)$ there are actually calculated over Ω (check this).

PROOF. By definition of v_{ε} ,

$$\begin{split} \int_{\Omega_{2\delta}} |v_{\varepsilon}(x+h) - v_{\varepsilon}(x)| \, \mathrm{d}x &= \int_{\Omega_{2\delta}} \left| \int_{\mathbf{R}^{N}} v(y) [\varphi_{\varepsilon}(x+h-y) - \varphi_{\varepsilon}(x-y)] \, \mathrm{d}y \right| \, \mathrm{d}x \\ &= \int_{\Omega_{2\delta}} \left| \int_{\mathbf{R}^{N}} \varphi_{\varepsilon}(y) [v(x+h-y) - v(x-y)] \, \mathrm{d}y \right| \, \mathrm{d}x \\ &\leq \int_{\mathbf{R}^{N}} \varphi_{\varepsilon}(y) \bigg[\int_{\Omega_{2\delta}} |v(x+h-y) - v(x-y)| \, \mathrm{d}x \bigg] \, \mathrm{d}y \\ &= \int_{\mathbf{R}^{N}} \varphi_{\varepsilon}(y) \bigg[\int_{\Omega_{2\delta}} |v(x+h-y) - v(x-y)| \, \mathrm{d}x \bigg] \, \mathrm{d}y \, . \end{split}$$

Here we use the standard notation

$$G - y = \{z \mid z + y \in G\}.$$

Recall that $\varphi_{\varepsilon}(y) = 0$ if $|y| \ge \varepsilon$. Therefore in last integral we may assume, as $\varepsilon < \delta$ by assumption,

$$\Omega_{2\delta} - y \subset \Omega_{2\delta - \varepsilon} \subset \Omega_{\delta} \,.$$

Hence the last integral above is bounded by

$$\int_{\mathbf{R}^N} \varphi_{\varepsilon}(y) \omega_{\delta}(|h|) \, \mathrm{d}y \le \omega_{\delta}(|h|) \, .$$

	٦.
	т

Let us define the sign function

$$\operatorname{sign}(s) = \begin{cases} 1, & s > 0, \\ 0, & s = 0, \\ -1, & s < 0. \end{cases}$$

Clearly, as $\varepsilon \to 0$,

$$v(x)[\operatorname{sign}(v(x))]_{\varepsilon} \to v(x)\operatorname{sign}(v(x)) = |v(x)|, \quad \text{a.e. } x \in \mathbb{R}^{N}.$$

Next lemma gives a measure of the speed of convergence in this limiting relation.

Lemma B.2. In the same assumptions of Lemma B.1, for all $0 < \varepsilon < \delta$,

$$\int_{\Omega_{\delta}} ||v(x)| - v(x)[\operatorname{sign}(v(x))]_{\varepsilon}| \, \mathrm{d}x \le 2\omega_{\delta}(\varepsilon) \,. \tag{B.4}$$

PROOF. For all $x, y \in \mathbf{R}^N$

$$\begin{aligned} ||v(x)| - v(x)\operatorname{sign}(v(y))| &= |[|v(x)| - |v(y)|] - [v(x) - v(y)]\operatorname{sign}(v(y))| \\ &\leq 2|v(x) - v(y)| \,. \end{aligned}$$

Thus

$$\int_{\Omega_{\delta}} ||v(x)| - v(x)[\operatorname{sign}(v(x))]_{\varepsilon}| \, \mathrm{d}x$$

$$= \int_{\Omega_{\delta}} \left| \int_{\mathbf{R}^{N}} \left[|v(x)| - v(x)\operatorname{sign}(v(y)) \right] \varphi_{\varepsilon}(x - y) \, \mathrm{d}y \right| \, \mathrm{d}x$$

$$\leq \int_{\Omega_{\delta}} \int_{\mathbf{R}^{N}} 2|v(x) - v(y)| \varphi_{\varepsilon}(x - y) \, \mathrm{d}y \, \mathrm{d}x = 2 \int_{\mathbf{R}^{N}} \int_{\Omega_{\delta}} |v(x) - v(x - y)| \, \mathrm{d}x \, \varphi_{\varepsilon}(y) \, \mathrm{d}y.$$

Since $\varphi_{\varepsilon}(y) = 0$ for $|y| \ge \varepsilon$, we may bound above last integral by

$$2\int_{\mathbf{R}^{N}}\int_{\Omega_{\delta}} |v(x) - v(x-y)| \, \mathrm{d}x \, \varphi_{\varepsilon}(y) \, \mathrm{d}y \leq 2\int_{\mathbf{R}^{N}} \omega_{\delta}(|y|) \varphi_{\varepsilon}(y) \, \mathrm{d}y \leq 2\omega_{\delta}(\varepsilon) \,.$$

B.2. The main estimate

Theorem B.3. Let $v \in L^{\infty}(Q_T)$, and assume that for all $1 > \delta > 0$, $|h| \leq \delta$ we have

$$\int_{\Omega_{\delta}} |v(x+h,t) - v(x,t)| \, \mathrm{d}x \le \omega_{\delta}(|h|), \qquad a.e. \ t \in (0,T). \tag{B.5}$$

Assume moreover that for all $g \in C_0^1(\Omega_{\delta})$, and a given $C_{\delta} > 0$,

$$\left| \int_{\Omega_{\delta}} g(x) [v(x,t+s) - v(x,t)] \, \mathrm{d}x \right| \le C_{\delta} s \left(\|g\|_{\infty} + \|\nabla g\|_{\infty} \right), \tag{B.6}$$

a.e. 0 < t < t + s < T. Then, we have a.e. 0 < t < t + s < T, and for all $0 < \varepsilon < \delta$,

$$\int_{\Omega_{\delta}} |v(x,t+s) - v(x,t)| \, \mathrm{d}x \le \gamma_{\delta} \left(\frac{s}{\varepsilon} + \varepsilon + \omega_{\delta}(\varepsilon)\right). \tag{B.7}$$

Here γ_{δ} depends on Ω , C_{δ} , $||v||_{\infty}$ and N.

PROOF. Choose in (B.6)

$$g(x) = \beta_{\varepsilon}(x) = \int_{\mathbf{R}^N} \beta(y) \varphi_{\varepsilon}(x-y) \,\mathrm{d}y,$$

where $0 < \varepsilon < \delta$ and

$$\beta(x) = \chi_{\Omega_{2\varepsilon+\delta}}(x)\operatorname{sign}(v(x,t+s) - v(x,t)).$$

Note that

$$|\nabla g| \le \frac{\gamma(N)}{\varepsilon};$$

moreover g(x) = 0 if $x \in \Omega_{\delta} - \Omega_{\varepsilon+\delta}$. Finally,

 $\beta_{\varepsilon}(x) = [\operatorname{sign}(v(x,t+s) - v(x,t))]_{\varepsilon}, \qquad x \in \Omega_{3\varepsilon+\delta}.$ Thus, exploiting assumption (B.6),

 $\left| \int_{\Omega_{3\varepsilon+\delta}} [v(x,t+s) - v(x,t)] \beta_{\varepsilon}(x) \, \mathrm{d}x \right| \leq \left| \int_{\Omega_{\delta}} [v(x,t+s) - v(x,t)] \beta_{\varepsilon}(x) \, \mathrm{d}x \right|$ $+ \left| \int_{\Omega_{\delta} - \Omega_{3\varepsilon+\delta}} [v(x,t+s) - v(x,t)] \beta_{\varepsilon}(x) \, \mathrm{d}x \right| \leq C_{\delta} \gamma(N) \frac{s}{\varepsilon} + \gamma(\Omega) \|v\|_{\infty} \varepsilon.$

Therefore

$$\begin{split} \int_{\Omega_{\delta}} |v(x,t+s) - v(x,t)| \, \mathrm{d}x &\leq \int_{\Omega_{3\varepsilon+\delta}} |v(x,t+s) - v(x,t)| \, \mathrm{d}x + \gamma(\Omega) \|v\|_{\infty} \varepsilon \\ &\leq \int_{\Omega_{3\varepsilon+\delta}} ||v(x,t+s) - v(x,t)| - [v(x,t+s) - v(x,t)] \beta_{\varepsilon}(x)| \, \mathrm{d}x + \\ &+ \left| \int_{\Omega_{3\varepsilon+\delta}} [v(x,t+s) - v(x,t)] \beta_{\varepsilon}(x) \, \mathrm{d}x \right| + \gamma(\Omega) \|v\|_{\infty} \varepsilon \leq 4\omega_{\delta}(\varepsilon) + \gamma \frac{s}{\varepsilon} + \gamma \varepsilon \,, \end{split}$$

where we used Lemma B.2. Indeed, it is easily checked, with the help of (B.5) that

$$\int_{\Omega_{\delta}} ||v(x+h,t+s) - v(x+h,t)| - |v(x,t+s) - v(x,t)|| \, \mathrm{d}x \le 2\omega_{\delta}(|h|) \,,$$

a.e. 0 < t < t + s < T, and for δ and h as above. The proof is concluded.

Remark B.4. It is clear that estimate (B.7) implies continuity in t of v in the $L^1(\Omega_{\delta})$ norm. In the optimal case when $\omega_{\delta}(|h|) = C_{\delta}|h|$, on choosing $\varepsilon = \sqrt{s}$ (for $s < \delta^2$) we obtain

$$\int_{\Omega_{\delta}} |v(x,t+s) - v(x,t)| \, \mathrm{d}x \le \gamma \sqrt{s} \,. \tag{B.8}$$

	-

APPENDIX C

The spaces $H^{m+\alpha,\frac{m+\alpha}{2}}(\overline{Q_T})$

We define here a class of Banach spaces of standard use in the regularity theory of parabolic equations (see [11]). Let $Q_T = \Omega \times (0,T)$, where $\Omega \subset \mathbf{R}^N$ is a bounded open set.

Fix the integer $m \geq 0$, and $\alpha \in (0,1)$. In the following we denote with $D_t^r D_x^s u$, for $r, s \in \mathbf{N}$, any derivative of u, taken r times with respect to the time variable, and s times with respect to space variables. For a given function $u: Q_T \to \mathbf{R}$ we introduce the quantities

$$\langle u \rangle_{x,Q_T}^{(\alpha)} = \sup_{(x,t),(x',t)\in Q_T} \frac{|u(x,t) - u(x',t)|}{|x - x'|^{\alpha}}, \quad \langle u \rangle_{x,Q_T}^{(m+\alpha)} = \sum_{2r+s=m} \langle D_t^r D_x^s u \rangle_{x,Q_T}^{(\alpha)},$$

$$\langle u \rangle_{t,Q_T}^{(\alpha/2)} = \sup_{(x,t),(x,t')\in Q_T} \frac{|u(x,t) - u(x,t')|}{|t - t'|^{\alpha/2}},$$

$$\langle u \rangle_{t,Q_T}^{(\frac{m+\alpha}{2})} = \sum_{2r+s=m-1,m} \langle D_t^r D_x^s u \rangle_{t,Q_T}^{\frac{(m+\alpha-2r-s)}{2}}.$$

The sums above (and below) are extended to all the derivatives $D_t^r D_x^s u$ with r, s as indicated. If m = 0, the only such function is u itself. Then we define the norm

$$|u|_{Q_T}^{(m+\alpha)} = \sum_{2r+s \le m} \|D_t^r D_x^s u\|_{\infty, Q_T} + \langle u \rangle_{x, Q_T}^{(m+\alpha)} + \langle u \rangle_{t, Q_T}^{(\frac{m+\alpha}{2})} + \langle u \rangle_$$

The Banach space of the functions u whose norm $|u|_{Q_T}^{(m+\alpha)}$ is finite is denoted by

$$H^{m+\alpha,\frac{m+\alpha}{2}}(\overline{Q_T})$$
.

C.1. Comments

The regularity 'in time' of $u \in H^{m+\alpha,\frac{m+\alpha}{2}}(\overline{Q_T})$ is 'a half' of the regularity 'in space', in a sense made precise by the definition itself. To illustrate this point, assume $s \in C([0,T])$, and regard s as a function defined over $\overline{Q_T}$. Let us make explicit the meaning of the statement

$$s \in H^{m+\alpha, \frac{m+\alpha}{2}}(\overline{Q_T}).$$

If m is even, m = 2k, this is equivalent to: $s \in C^k([0,T])$, and

$$\sup_{0 < t, t' < T} \frac{|s^{(k)}(t) - s^{(k)}(t')|}{|t - t'|^{\frac{\alpha}{2}}} < \infty \,.$$

If instead m is odd, m = 2k - 1, we have: $s \in C^{k-1}([0,T])$, and

$$\sup_{0 < t, t' < T} \frac{|s^{(k-1)}(t) - s^{(k-1)}(t')|}{|t - t'|^{\frac{1+\alpha}{2}}} < \infty.$$

Clearly, for all $m \ge 0$,

$$\dot{s} \in H^{m+\alpha,\frac{m+\alpha}{2}}(\overline{Q_T}) \Longrightarrow s \in H^{2+m+\alpha,\frac{2+m+\alpha}{2}}(\overline{Q_T}).$$
(C.1)
APPENDIX D

Solution of a simple variational inequality

D.1. A variational inequality

We are interested here in solutions of the following problem: (P): Find a function $\psi \in W_1^2(0,T)$ such that:

• For a given $\psi_0 \in [0, 1]$

$$\psi(0) = \psi_0 \,. \tag{D.1}$$

• The function ψ is bounded so as

$$0 \le \psi(t) \le 1$$
, $0 \le t \le T$. (D.2)

- For a given $f \in L^2(0,T)$ $\psi'(t) = f(t)$, for a.e. $0 \le t \le T$ such that $0 < \psi(t) < 1$. (D.3)
- For a.e. t

$$\psi(t) = 0 \quad \Rightarrow \quad f(t) \le 0; \qquad \psi(t) = 1 \quad \Rightarrow \quad f(t) \ge 0.$$
 (D.4)

Remark D.1. Problem (P) is a mathematical model for an evolution phenomenon with unilateral constraints; in other words, ψ obeys the o.d.e. in (D.3) as far as this is possible without trespassing the bounds in (D.2).

Remark D.2. Clearly the restriction $0 < \psi(t) < 1$ in requirement (D.3) is necessary for the problem to have any meaning. For example, assume $\psi_0 = 0$ and f(t) = -1 for all t. A solution to (P) is given by the constant $\psi = 0$. It is easy to see that this is the only function in $W_2^1(0,T)$ satisfying (P). Therefore the restriction $0 < \psi(t) < 1$ can not be dropped in general if existence of solutions is to be preserved.

A deeper feature of our formulation of problem (P) must be pointed out: assume $\psi_0 = 0$ and f(t) = 1 for all t. Then a solution is given by $\psi(t) = t$, at least up to time T = 1. But the constant function $\psi = 0$ again satisfies (D.1)–(D.3), though its behaviour is not in the spirit of Remark D.1, and in fact (D.4) is not fulfilled.

Remark D.3. One can attempt to construct a solution to (P) by means of the following intuitive procedure: say $0 < \psi_0 < 1$; then define ψ as the solution to the Cauchy problem for the o.d.e. $\psi' = f$ with initial data ψ_0 , in the maximal interval $(0, t_0)$ where $0 < \psi < 1$. Say e.g., $\psi(t_0) = 0$. Let $t_1 > t_0$ be the first time greater than t_0 such that $f \ge 0$ (but f is not identically vanishing) in $(t_1, t_1 + \delta)$ for some $\delta > 0$. Then set $\psi(t) = 0$ for $t_0 < t < t_1$ and define again ψ as the solution to $\psi' = f$, $\psi(t_1) = 0$ in $(t_1, t_1 + \delta)$, so that we are essentially back to the case of initial data in the open interval (0, 1), since $\psi(t_1 + \delta) > 0$. Unfortunately this approach is too naive: for example the first time t_1 with the properties above needs not exist, even if f > 0 in a set of positive measure for $t > t_0$.

DANIELE ANDREUCCI

In order to circumvent the difficulties we described in Remark D.3, we take in next Section a different approach, based on a technique known as *penalization*.

D.2. Existence of solutions to (P)

Define a sequence $f_n \in C([0,T])$ such that

$$f_n \to f$$
, in $L^2(0,T)$, and a.e. in $(0,T)$, as $n \to \infty$. (D.5)

Also define the sequence of functions

$$\eta_n(s) = \begin{cases} ns, & s < 0, \\ 0, & 0 \le s \le 1, \\ n(s-1), & s > 1. \end{cases}$$
(D.6)

In Figure D.1 we sketch the graphs of η_n and of their limit η . Notice that we do not give here a rigorous definition of the limiting relation $\eta_n \to \eta$. In



FIGURE D.1. The approximating functions η_n and their limit η .

the same spirit, we just appeal to Figure D.1 to support the statement that η can be regarded as the inverse of the Heaviside graph

$$H(s) = \begin{cases} 0, & s < 0, \\ [0,1], & s = 0, \\ 1, & s > 0. \end{cases}$$

In the following we denote by $\psi_n \in C^1([0,T])$ the solution to the Cauchy problem

$$\frac{\mathrm{d}\psi_n}{\mathrm{d}t} + \eta_n(\psi_n) = f_n \,, \tag{D.7}$$

$$\psi_n(0) = \psi_0 \,. \tag{D.8}$$

Such a solution exists by virtue of standard results on classical o.d.e.; in general one may have $\psi_n(t) \notin [0, 1]$ for some t.

Lemma D.4. Perhaps by extracting a subsequence, which we still denote by $\{\psi_n\}$, we may assume

$$\psi_n \to \psi$$
, in $L^2(0,T)$ and a.e. in $(0,T)$; (D.9)

$$\frac{\mathrm{d}\psi_n}{\mathrm{d}t} \to \frac{\mathrm{d}\psi}{\mathrm{d}t}, \qquad weakly \ in \ L^2(0,T), \tag{D.10}$$

for a suitable $\psi \in W_2^1(0,T)$. Moreover

$$0 \le \psi \le 1$$
, *a.e.* in $(0, T)$. (D.11)

PROOF. On multiplying (D.7) by ψ'_n and integrating over (0, t) we obtain

$$\begin{split} \int_{0}^{t} \left| \frac{\mathrm{d}\psi_{n}}{\mathrm{d}\tau}(\tau) \right|^{2} \,\mathrm{d}\tau + \int_{0}^{t} \eta_{n}(\psi_{n}(\tau)) \frac{\mathrm{d}\psi_{n}}{\mathrm{d}\tau}(\tau) \,\mathrm{d}\tau = \int_{0}^{t} f_{n}(\tau) \frac{\mathrm{d}\psi_{n}}{\mathrm{d}\tau}(\tau) \,\mathrm{d}\tau \\ & \leq \frac{1}{2} \int_{0}^{t} f_{n}(\tau)^{2} \,\mathrm{d}\tau + \frac{1}{2} \int_{0}^{t} \left| \frac{\mathrm{d}\psi_{n}}{\mathrm{d}\tau}(\tau) \right|^{2} \,\mathrm{d}\tau \,, \end{split}$$

whence

$$\int_{0}^{t} \left| \frac{\mathrm{d}\psi_{n}}{\mathrm{d}\tau}(\tau) \right|^{2} \mathrm{d}\tau + 2 \int_{\psi_{0}}^{\psi_{n}(t)} \eta_{n}(s) \,\mathrm{d}s \leq \gamma \int_{0}^{t} f(\tau)^{2} \,\mathrm{d}\tau \leq \gamma \|f\|_{2,(0,T)}^{2}. \quad (\mathrm{D}.12)$$

Since for every $t \in (0, T]$, by Hölder inequality,

$$|\psi_n(t)| \le \psi_0 + \int_0^t \left| \frac{\mathrm{d}\psi_n}{\mathrm{d}\tau}(\tau) \right| \,\mathrm{d}\tau \le \psi_0 + \sqrt{t} \left\| \frac{\mathrm{d}\psi_n}{\mathrm{d}\tau} \right\|_{2,(0,t)} \le \gamma$$

standard results imply both (D.9) and (D.10). Assume next that $\psi_n(t) > 1$ for given $t \in [0, T]$ and $n \ge 1$. By (D.12) we infer

$$\gamma \ge \int_{\psi_0}^{\psi_n(t)} \eta_n(s) \, \mathrm{d}s = \int_{1}^{\psi_n(t)} \eta_n(s) \, \mathrm{d}s = \frac{n}{2} (\psi_n(t) - 1)^2 \, .$$

From this estimate and a similar one valid for the case $\psi_n(t) < 0$ we get

$$-\frac{\gamma}{\sqrt{n}} \le \psi_n(t) \le 1 + \frac{\gamma}{\sqrt{n}}, \qquad 0 \le t \le T.$$
 (D.13)

This proves (D.11).

Lemma D.5. Perhaps by extracting a subsequence, which we still denote by $\{\psi_n\}$, we may assume as $n \to \infty$

$$\eta_n(\psi_n) \to \xi = -\xi_0 + \xi_1, \qquad \text{weakly in } L^2(0,T),$$
 (D.14)

where both ξ_i , i = 0, 1 are nonnegative functions, such that $\xi_i = 0$ a.e. outside of the set $\{\psi = i\}$, for i = 0 and i = 1 respectively.

PROOF. On multiplying (D.7) by $\eta_n(\psi_n)$ and integrating over (0,t) we obtain

$$\int_{0}^{t} \eta_{n}(\psi_{n}(\tau)) \frac{\mathrm{d}\psi_{n}}{\mathrm{d}\tau}(\tau) \,\mathrm{d}\tau + \int_{0}^{t} |\eta_{n}(\psi_{n}(\tau))|^{2} \,\mathrm{d}\tau = \int_{0}^{t} f_{n}(\tau)\eta_{n}(\psi_{n}(\tau)) \,\mathrm{d}\tau$$
$$\leq \frac{1}{2} \int_{0}^{t} f_{n}(\tau)^{2} \,\mathrm{d}\tau + \frac{1}{2} \int_{0}^{t} |\eta_{n}(\psi_{n}(\tau))|^{2} \,\mathrm{d}\tau$$

whence

$$2 \int_{\psi_0}^{\psi_n(t)} \eta_n(s) \,\mathrm{d}s + \int_0^t |\eta_n(\psi_n(\tau))|^2 \,\mathrm{d}\tau \le \gamma \int_0^t f(\tau)^2 \,\mathrm{d}\tau \le \gamma ||f||_{2,(0,T)}^2.$$
(D.15)

Thus the weak convergence $\eta_n(\psi_n) \to \xi$ for some $\xi \in L^2(0,T)$ follows from standard results and the uniform L^2 estimate in (D.15). Since by linearity we may define ξ_i as the weak limits in

$$-\eta_n(\psi_n)\chi_{\{\psi_n<0\}} \to \xi_0 , \qquad \eta_n(\psi_n)\chi_{\{\psi_n>1\}} \to \xi_1 ,$$

it is only left to prove the assertion about the support of ξ_i . For example we calculate

$$0 \le \int_{0}^{T} \xi_{0} \psi \, \mathrm{d}t = \lim_{n \to \infty} \int_{0}^{T} (-\eta_{n}(\psi_{n})\chi_{\{\psi_{n} < 0\}})\psi_{n} \, \mathrm{d}t \le 0 \,,$$

proving that $\xi_0 \psi = 0$ a.e. in (0, T). In the same way we prove that $\xi_1(1-\psi) = 0$ a.e. in (0, T).

Proposition D.6. The function ψ obtained in Lemma D.4, D.5 solves problem (P).

PROOF. We show that (D.3) is in force. Observe that as $n \to \infty$

$$\int_{0}^{T} \left| \frac{\mathrm{d}\psi_{n}}{\mathrm{d}t} - f_{n} \right| \chi_{\{\psi \in (0,1)\}} \,\mathrm{d}t = \int_{0}^{T} |\eta_{n}(\psi_{n})| \chi_{\{\psi \in (0,1)\}} \,\mathrm{d}t \to 0 \,,$$

invoking the uniform integrability in (D.15), and by virtue of

$$\eta_n(\psi_n(t)) \to 0$$
, a.e. t such that $0 < \psi(t) < 1$,

which follows from the definition of η_n . Then, perhaps by extracting a subsequence again, we may assume as $n \to \infty$

$$\frac{\mathrm{d}\psi_n}{\mathrm{d}t} \to f\,, \qquad \text{a.e. in } \{0 < \psi < 1\},$$

implying $\psi' = f$ in $\{0 < \psi < 1\}$.

Condition (D.4) follows from Exercise D.8, and from Lemma D.5. $\hfill \Box$

Remark D.7. From condition (D.3) we readily infer

$$\left\| \frac{\mathrm{d}\psi}{\mathrm{d}t} \right\|_{2,(0,t)}, \|\xi\|_{2,(0,t)} \le \|f\|_{2,(0,t)}, \qquad 0 < t < T.$$
(D.16)

D.2.1. Exercises.

Exercise D.8. Prove that $\xi_0 = -f$ a.e. in $\{\xi_0 \neq 0\}$, and that $\xi_1 = f$ a.e. in $\{\xi_1 \neq 0\}$.

D.3. Variational formulation

Recalling the results of Section D.2, it is quite natural to rewrite formally problem (P) as

$$\frac{\mathrm{d}\psi}{\mathrm{d}t} + \eta(\psi) \ni f \,, \tag{D.17}$$

$$\psi(0) = \psi_0 \in [0, 1]. \tag{D.18}$$

Definition D.9. A function $\psi \in W_1^2(0,T)$ is a solution to the variational inequality (D.17), (D.18) if $0 \le \psi \le 1$, (D.18) is satisfied in the standard pointwise sense (remember that $W_1^2(0,T) \subset C([0,T])$), and

$$\int_{0}^{T} \left(\frac{\mathrm{d}\psi}{\mathrm{d}t} - f\right) (\varphi - \psi) \,\mathrm{d}t \ge 0 \tag{D.19}$$

for all $\varphi \in L^{\infty}(0,T), 0 \leq \varphi \leq 1$.

Theorem D.10. The function ψ obtained in Lemma D.4, D.5 is the unique solution to problem (P) and to the variational inequality.

PROOF. A) Assume a function $\hat{\psi}$ solves (P), and let φ be as in Definition D.9. Thus obviously

$$\left(\frac{\mathrm{d}\hat{\psi}}{\mathrm{d}t} - f\right)(\varphi - \hat{\psi}) = 0, \quad \text{if } 0 < \hat{\psi} < 1.$$

Moreover by condition (D.4) we have

$$\left(\frac{\mathrm{d}\psi}{\mathrm{d}t} - f\right)(\varphi - \hat{\psi}) = -f(\varphi - i) \ge 0, \qquad \text{in } \{\hat{\psi} = i\}, \, i = 0, \, 1.$$

B) To complete the proof of the Theorem, in view of the existence result of Proposition D.6, we need only prove uniqueness of solutions for the variational inequality.

Let ψ_a , ψ_b be two such solutions. We add (D.19) written for ψ_a with $\varphi = \psi_b$ to (D.19) written for ψ_b with $\varphi = \psi_a$, obtaining

$$\int_{0}^{T} \left(\frac{\mathrm{d}\psi_{a}}{\mathrm{d}t} - \frac{\mathrm{d}\psi_{b}}{\mathrm{d}t} \right) (\psi_{b} - \psi_{a}) \, \mathrm{d}t \ge 0 \,,$$

yielding upon integration

$$-(\psi_b(T) - \psi_a(T))^2 \ge 0,$$

i.e., $\psi_b(T) = \psi_a(T)$. Since T in this connection can be replaced with any $t \in (0, T)$, uniqueness is proven.

•

Proposition D.11. Let ψ_i i = 1, 2 be two solutions of the variational inequality corresponding respectively to $f = f_i$, $\psi_0 = \psi_{0i}$. Then for all 0 < t < T

$$(\psi_1(t) - \psi_2(t))^2 \le e^t (\psi_{01} - \psi_{02})^2 + (e^t - 1) \|f_1 - f_2\|_{2,(0,t)}^2.$$
(D.20)

PROOF. We add (D.19) written for ψ_1 with $\varphi = \psi_2$ to (D.19) written for ψ_2 with $\varphi = \psi_1$, obtaining

$$\int_{0}^{t} \left(\frac{\mathrm{d}\psi_{1}}{\mathrm{d}\tau} - \frac{\mathrm{d}\psi_{2}}{\mathrm{d}\tau}\right) (\psi_{1} - \psi_{2}) \,\mathrm{d}\tau \leq \int_{0}^{t} (f_{1} - f_{2})(\psi_{1} - \psi_{2}) \,\mathrm{d}\tau \,. \tag{D.21}$$

By integration of the left hand side, and after an application of Cauchy-Schwarz inequality we get

$$\frac{1}{2}(\psi_1(t) - \psi_2(t))^2 - \frac{1}{2}(\psi_{01} - \psi_{02})^2 \le \frac{1}{2}\int_0^t (f_1 - f_2)^2 \,\mathrm{d}\tau + \frac{1}{2}\int_0^t (\psi_1 - \psi_2)^2 \,\mathrm{d}\tau \,,$$

whence (D.20) follows by virtue of Gronwall's lemma.

D.3.1. Dependence on the space variable. In view of the application of Chapter 2, we are interested in the case where problem (P) or equivalently the variational inequality are parametrized by a space variable $x \in \Omega$. Thus $f \in L^2(Q_T), \psi_0 \in L^{\infty}(\Omega), 0 \leq \psi_0 \leq 1$, and in Definition D.9 we assume $\psi \in W_1^2(0,T; L^2(\Omega))$; also, (D.19) is assumed to be valid for a.e. $x \in \Omega$. All the proofs in this Appendix stay essentially unchanged. We only need one

For the proofs in this Appendix stay essentially unchanged. We only need one more ingredient to prove compactness in the $L^2(Q_T)$ sense of the sequence ψ_n , i.e., an equicontinuity estimate in the $L^2(Q_T)$ norm with respect to the space variables, the case of the time variable being handled as above. This is provided by the following argument. Denote for any $\delta > 0$

$$\Omega_{\delta} = \{ x \in \Omega \mid \operatorname{dist}(x, \partial \Omega) > \delta \},\$$
$$g^{h}(x, t) = g(x + h, t),\$$

where $h \in \mathbf{R}^N$ is an arbitrary vector such that $|h| < \delta$, and g is any function defined in Q_T . We have

$$\begin{split} \int_{\Omega_{\delta}(t)} \left| \psi_{n}^{h} - \psi_{n} \right|^{2} \mathrm{d}x &= \int_{\Omega_{\delta}} \left| \psi_{0}^{h} - \psi_{0} \right|^{2} \mathrm{d}x \\ &+ 2 \int_{0}^{t} \int_{\Omega_{\delta}} (\psi_{n}^{h} - \psi_{n}) \Big(\frac{\partial \psi_{n}^{h}}{\partial \tau} - \frac{\partial \psi_{n}}{\partial \tau} \Big) \mathrm{d}x \, \mathrm{d}\tau \,. \end{split}$$

Since η_n is non-decreasing, the last integral above can be majorised by

$$(\psi_n^h - \psi_n) \Big(\frac{\partial \psi_n^h}{\partial \tau} - \frac{\partial \psi_n}{\partial \tau} \Big) \le (\psi_n^h - \psi_n) \Big(\frac{\partial \psi_n^h}{\partial \tau} - \frac{\partial \psi_n}{\partial \tau} \Big)$$

+ $[\eta_n(\psi_n^h) - \eta_n(\psi_n)](\psi_n^h - \psi_n) = (f_n^h - f_n)(\psi_n^h - \psi_n) .$

Collecting the estimates above we get

$$\begin{split} \int\limits_{\Omega_{\delta}(t)} \left| \psi_{n}^{h} - \psi_{n} \right|^{2} \, \mathrm{d}x &\leq \int\limits_{\Omega_{\delta}} \left| \psi_{0}^{h} - \psi_{0} \right|^{2} \, \mathrm{d}x \\ &+ \int\limits_{0}^{t} \int\limits_{\Omega_{\delta}} |f_{n}^{h} - f_{n}|^{2} \, \mathrm{d}x \, \mathrm{d}\tau + \int\limits_{0}^{t} \int\limits_{\Omega_{\delta}} \left| \psi_{n}^{h} - \psi_{n} \right|^{2} \, \mathrm{d}x \, \mathrm{d}\tau \,, \end{split}$$

so that the required estimate

$$\begin{split} \|\psi_n^h - \psi_n\|_{2,\Omega_{\delta}(t)} &\leq \gamma \|\psi_0^h - \psi_0\|_{2,\Omega_{\delta}} + \gamma \|f^h - f\|_{2,\Omega_{\delta} \times (0,T)} \\ \text{follows from a standard application of Gronwall's lemma and from the convergence of } f_n \text{ to } f \text{ in } L^2(Q_T). \end{split}$$

APPENDIX E

Symbols used in text

$\ f\ _{p,D}$	norm of f in $L^p(D)$, $1 \le p \le \infty$; often the indication of the domain
	is omitted, when superfluous.
$\Omega(t)$	the set $\{(x,t) \mid x \in \Omega\}$.
$ \Omega $	the Lebesgue measure of the set Ω .
$\operatorname{dist}(A, B)$	standard Euclidean distance between A and B .
$B_r(x)$	ball with center x and radius r .
$x \rightarrow s_0 +$	x goes to s_0 from the right.
$x \to s_0 -$	x goes to s_0 from the left.
$f(s_0+)$	denotes the limit of $f(x)$ as $x \to s_0 +$.
$f(s_0-)$	denotes the limit of $f(x)$ as $x \to s_0 -$.
s_+	positive part of $s \in \mathbf{R}$, $s_+ = \max(s, 0)$.
<i>S</i> _	negative part of $s \in \mathbf{R}$, $s_{-} = \max(-s, 0)$.
δ_{ij}	Kronecker's symbol: $\delta_{ij} = 1$ if $i = j$, $\delta_{ij} = 0$ if $i \neq j$.
∇f	spatial gradient of the function $f(x,t)$: $\nabla f = (\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_N}).$
$\mathbf{e_k}$	k-th unit vector of the standard basis in \mathbf{R}^N .
C(A)	class of continuous functions in A. The same as $C^{0}(A)$.
$C^n(A)$	class of functions, continuous in A together with
	their derivatives up to order n .
$C_0^n(A)$	class of the functions in $C^{n}(A)$, whose support is
	compact and contained in A .
$C^{2,1}(A)$	class of functions f , such that f , $\frac{\partial f}{\partial t}$, $\frac{\partial f}{\partial x_i}$, and $\frac{\partial^2 f}{\partial x_i \partial x_j}$
2	are continuous in A for all $i, j = 1,, N$.
$W_{1}^{2}(G)$	Sobolev space of functions in $L^2(G)$ whose first derivatives are in $L^2(G)$.
$W_{2,1}^2(G)$	Sobolev space of functions $f \in W_1^2(G)$ whose spatial derivatives
	$\frac{\partial^2 f}{\partial x_i \partial x_j}$ are in $L^2(G)$ for all $i, j = 1, \dots, N$.
PDE	Partial Differential Equation/Equations.

Bibliography

- D. Andreucci. Behaviour of mushy regions under the action of a volumetric heat source. Math. Methods in the Applied Sciences, 16:35–47, 1993.
- [2] O. Arena. *Partial Differential Equations of Parabolic Type*. Notes of a course held at the University of Florence (Italy). Unpublished, 1983.
- [3] J. R. Cannon. The One-Dimensional Heat Equation, volume 23 of Encyclopedia of Mathematics and Its Applications. Addison-Wesley, Menlo Park, CA U.S.A., 1984.
- [4] E. DiBenedetto. Continuity of weak solutions to certain singular parabolic equations. Ann. Mat. Pura Appl., 130:131–176, 1982.
- [5] A. Fasano. Free Boundary problems and Their Applications. Notes of a course held at SASIAM, Bari (Italy). 1989.
- [6] A. Fasano and M. Primicerio. New results on some classical parabolic free-boundary problems. *Quart. Appl. Math.*, 38:439–460, 1981.
- [7] A. Friedman. Remarks on the maximum principle for parabolic equations and its applications. *Pacific J. Math.*, 8:201–211, 1958.
- [8] I. G. Götz and B. B. Zaltzman. Nonincrease of mushy regions in a nonhomogeneous Stefan problem. *Quart. Appl. Math.*, 49:741–746, 1991.
- [9] S. L. Kamenomostskaya. On Stefan problem. Mat. Sbornik, 53:489–514, 1961. (In Russian).
- [10] S. N. Kruzhkov. First-order quasilinear equations in several independent variables. Mat. Sb. (n.s.), 81:228–255, 1970. Engl. Transl.: Math. USSR - Sb 10 (1970), 217-234.
- [11] O. A. Ladyzhenskaja, V. A. Solonnikov, and N. N. Ural'ceva. Linear and Quasilinear Equations of Parabolic Type, volume 23 of Translations of Mathematical Monographs. American Mathematical Society, Providence, RI, 1968.
- [12] A. M. Meirmanov. The Stefan Problem, volume 3 of de Gruyter Expositions in Mathematics. Walter de Gruyter, Berlin, Germany, 1992.
- [13] L. Nirenberg. A strong maximum principle for parabolic equations. Comm. Pure Appl. Math., 6:167–177, 1953.
- [14] O. A. Oleinik. A method of solution of the general Stefan problem. Soviet Math. Dokl., 1:1350–1354, 1960.
- [15] M. Primicerio. Mushy regions in phase-change problems. In K. H. Hoffmann and R. Gorenflo, editors, Applied Nonlinear Functional Analysis: Variational Methods and Ill-Posed Problems, pages 251–269, Frankfurt, Germany, 1983. Verlag Peter Lang.
- [16] D. G. Schaeffer. A new proof of the infinite differentiability of the free boundary in the Stefan problem. J. Diff. Equations, 20:266–269, 1976.
- [17] J. Stefan. Über die Theorie der Eisbildung. Monatshefte Mat. Phys., 1:1-6, 1890.
- [18] D. Tarzia. A bibliography on moving-free boundary problems for the heat-diffusion equation. The Stefan and related problems. *MAT-Serie A*, 2, 2000.
- http://web.austral.edu.ar/cienciasEmpresariales-investigacion-mat-A-02.asp. [19] R. Viborni. On properties of solutions of some boundary value problems for equations
- of parabolic type. Doklady Akad. Nauk SSSR (n.s.), 117:563–565, 1957. (In Russian). [20] A. Visintin. Models of Phase Transition, volume 28 of Progress in Nonlinear Differ-